arXiv:1203.4433v2 [math.DG] 6 Nov 2012

# THE COTTON TENSOR AND THE RICCI FLOW

CARLO MANTEGAZZA, SAMUELE MONGODI, AND MICHELE RIMOLDI

ABSTRACT. We compute the evolution equation of the Cotton tensor under the Ricci flow of a Riemannian manifold, with particular attention to the three dimensional case, and we discuss some applications.

## CONTENTS

1. Preliminaries 1
2. The Evolution Equation of the Cotton Tensor in 3D 2
3. Three–Dimensional Gradient Ricci Solitons 10
4. The Evolution Equation of the Cotton Tensor in any Dimension 13
References 23

## 1. PRELIMINARIES

The Riemann curvature operator of a Riemannian manifold $(M^n, g)$ is defined, as in [5], by

$$\mathrm{Riem}(X,Y)Z = \nabla_Y\nabla_X Z - \nabla_X\nabla_Y Z + \nabla_{[X,Y]}Z\,.$$

In a local coordinate system the components of the $(3,1)$–Riemann curvature tensor are given by $\mathrm{R}^l_{ijk}\frac{\partial}{\partial x^l} = \mathrm{Riem}\big(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\big)\frac{\partial}{\partial x^k}$ and we denote by $\mathrm{R}_{ijkl} = g_{lm}\mathrm{R}^m_{ijk}$ its $(4,0)$–version.

With the previous choice, for the sphere $\mathbb{S}^n$ we have $\mathrm{Riem}(v,w,v,w) = \mathrm{R}_{abcd}v^iw^jv^kw^l > 0$.

*In all the paper the Einstein convention of summing over the repeated indices will be adopted.*

The Ricci tensor is obtained by the contraction $\mathrm{R}_{ik} = g^{jl}\mathrm{R}_{ijkl}$ and $\mathrm{R} = g^{ik}\mathrm{R}_{ik}$ will denote the scalar curvature.

We recall the interchange of derivative formula,

$$\nabla^2_{ij}\omega_k - \nabla^2_{ji}\omega_k = \mathrm{R}_{ijkp}g^{pq}\omega_q\,,$$

and Schur's lemma, which follows by the second Bianchi identity,

$$2g^{pq}\nabla_p\mathrm{R}_{qi} = \nabla_i\mathrm{R}\,.$$

They both will be used extensively in the computations that follows.

The so called Weyl tensor is then defined by the following decomposition formula (see [5, Chapter 3, Section K]) in dimension $n \geq 3$,

$$\mathrm{R}_{ijkl} = \frac{1}{n-2}(\mathrm{R}_{ik}g_{jl} - \mathrm{R}_{il}g_{jk} + \mathrm{R}_{jl}g_{ik} - \mathrm{R}_{jk}g_{il}) - \frac{\mathrm{R}}{(n-1)(n-2)}(g_{ik}g_{jl} - g_{il}g_{jk}) + \mathrm{W}_{ijkl}\,. \tag{1.1}$$

The Weyl tensor satisfies all the symmetries of the curvature tensor, moreover, all its traces with the metric are zero, as it can be easily seen by the above formula.
In dimension three W is identically zero for every Riemannian manifold. It becomes relevant instead when $n \geq 4$ since its vanishing is a condition equivalent for $(M^n, g)$ to be *locally conformally flat*, that is, around every point $p \in M^n$ there is a conformal deformation $\widetilde{g}_{ij} = e^f g_{ij}$ of the original metric $g$, such that the new metric is flat, namely, the Riemann tensor associated to $\widetilde{g}$ is zero in $U_p$ (here $f : U_p \to \mathbb{R}$ is a smooth function defined in a open neighborhood $U_p$ of $p$).
In dimension $n = 3$, instead, locally conformally flatness is equivalent to the vanishing of the following *Cotton tensor*

$$\mathrm{C}_{ijk} = \nabla_k\mathrm{R}_{ij} - \nabla_j\mathrm{R}_{ik} - \frac{1}{2(n-1)}\big(\nabla_k\mathrm{R}g_{ij} - \nabla_j\mathrm{R}g_{ik}\big)\,, \tag{1.2}$$

---

*Date*: February 27, 2013.

which expresses the fact that the *Schouten* tensor

$$\mathrm{S}_{ij} = \mathrm{R}_{ij} - \frac{\mathrm{R}g_{ij}}{2(n-1)}$$

is a *Codazzi* tensor (see [1, Chapter 16, Section C]), that is, a symmetric bilinear form $\mathrm{T}_{ij}$ such that $\nabla_k \mathrm{T}_{ij} = \nabla_i \mathrm{T}_{kj}$.

By means of the second Bianchi identity, one can easily get (see [1]) that

$$\nabla^l \mathrm{W}_{lijk} = -\frac{n-3}{n-2}\mathrm{C}_{ijk}\,. \tag{1.3}$$

Hence, when $n \geq 4$, if we assume that the manifold is locally conformally flat (that is, $\mathrm{W} = 0$), the Cotton tensor is identically zero also in this case, but this is only a necessary condition.

By direct computation, we can see that the tensor $\mathrm{C}_{ijk}$ satisfies the following symmetries

$$\mathrm{C}_{ijk} = -\mathrm{C}_{ikj}, \qquad \mathrm{C}_{ijk} + \mathrm{C}_{jki} + \mathrm{C}_{kij} = 0\,, \tag{1.4}$$

moreover it is trace–free in any two indices,

$$g^{ij}\mathrm{C}_{ijk} = g^{ik}\mathrm{C}_{ijk} = g^{jk}\mathrm{C}_{ijk} = 0\,, \tag{1.5}$$

by its skew–symmetry and Schur's lemma.

We suppose now that $(M^n, g(t))$ is a Ricci flow in some time interval, that is, the time–dependent metric $g(t)$ satisfies

$$\frac{\partial}{\partial t} g_{ij} = -2\mathrm{R}_{ij}.$$

We have then the following evolution equations for the Christoffel symbols, the Ricci tensor and the scalar curvature (see for instance [6]),

$$\begin{aligned}
\frac{\partial}{\partial t}\Gamma^k_{ij} &= -g^{ks}\nabla_i \mathrm{R}_{js} - g^{ks}\nabla_j \mathrm{R}_{is} + g^{ks}\nabla_s \mathrm{R}_{ij} \\
\frac{\partial}{\partial t}\mathrm{R}_{ij} &= \Delta \mathrm{R}_{ij} - 2\mathrm{R}^{kl}\mathrm{R}_{kijl} - 2g^{pq}\mathrm{R}_{ip}\mathrm{R}_{jq} \\
\frac{\partial}{\partial t}\mathrm{R} &= \Delta \mathrm{R} + 2|\mathrm{Ric}|^2\,.
\end{aligned} \tag{1.6}$$

*All the computations which follow will be done in a fixed local frame, not in a moving frame.*

**Acknowledgments.** *The second and third authors are partially supported by the Italian FIRB Ideas "Analysis and Beyond".*

*Note.* We remark that Huai-Dong Cao also, indipendently by us, worked out the computation of the evolution of the Cotton tensor in dimension three, in an unpublished note.

## 2. THE EVOLUTION EQUATION OF THE COTTON TENSOR IN 3D

The goal of this section is to compute the evolution equation under the Ricci flow of the Cotton tensor $\mathrm{C}_{ijk}$ in dimension three (see [4] for the evolution of the Weyl tensor), the general computation in any dimension is postponed to section 4.

In the special threedimensional case we have,

$$\mathrm{R}_{ijkl} = \mathrm{R}_{ik}g_{jl} - \mathrm{R}_{il}g_{jk} + \mathrm{R}_{jl}g_{ik} - \mathrm{R}_{jk}g_{il} - \frac{\mathrm{R}}{2}(g_{ik}g_{jl} - g_{il}g_{jk})\,, \tag{2.1}$$

$$\mathrm{C}_{ijk} = \nabla_k \mathrm{R}_{ij} - \nabla_j \mathrm{R}_{ik} - \frac{1}{4}\big(\nabla_k \mathrm{R} g_{ij} - \nabla_j \mathrm{R} g_{ik}\big)\,, \tag{2.2}$$

hence, the evolution equations (1.6) become

$$
\begin{aligned}
\frac{\partial}{\partial t}\Gamma^k_{ij} &= -\, g^{ks}\nabla_i \mathrm{R}_{js} - g^{ks}\nabla_j \mathrm{R}_{is} + g^{ks}\nabla_s \mathrm{R}_{ij}\\
\frac{\partial}{\partial t}\mathrm{R}_{ij} &= \Delta \mathrm{R}_{ij} - 6g^{pq}\mathrm{R}_{ip}\mathrm{R}_{jq} + 3\mathrm{R}\mathrm{R}_{ij} + 2|\mathrm{Ric}|^2 g_{ij} - \mathrm{R}^2 g_{ij}\\
\frac{\partial}{\partial t}\mathrm{R} &= \Delta \mathrm{R} + 2|\mathrm{Ric}|^2\,.
\end{aligned}
$$

From these formulas we can compute the evolution equations of the derivatives of the curvatures assuming, from now on, to be in normal coordinates,

$$
\begin{aligned}
\frac{\partial}{\partial t}\nabla_l \mathrm{R} \quad &= \quad \nabla_l \Delta \mathrm{R} + 2\nabla_l |\mathrm{Ric}|^2\,,\\
\frac{\partial}{\partial t}\nabla_s \mathrm{R}_{ij} \quad &= \quad \nabla_s \Delta \mathrm{R}_{ij} - 6\nabla_s \mathrm{R}_{ip}\mathrm{R}_{jp} - 6\mathrm{R}_{ip}\nabla_s \mathrm{R}_{jp} + 3\nabla_s \mathrm{R}\mathrm{R}_{ij} + 3\mathrm{R}\nabla_s \mathrm{R}_{ij}\\
&\qquad +2\nabla_s|\mathrm{Ric}|^2 g_{ij} - \nabla_s \mathrm{R}^2 g_{ij}\\
&\qquad +(\nabla_i \mathrm{R}_{sp} + \nabla_s \mathrm{R}_{ip} - \nabla_p \mathrm{R}_{is})\mathrm{R}_{jp}\\
&\qquad +(\nabla_j \mathrm{R}_{sp} + \nabla_s \mathrm{R}_{jp} - \nabla_p \mathrm{R}_{js})\mathrm{R}_{ip}\\
&= \quad \nabla_s \Delta \mathrm{R}_{ij} - 5\nabla_s \mathrm{R}_{ip}\mathrm{R}_{jp} - 5\mathrm{R}_{ip}\nabla_s \mathrm{R}_{jp} + 3\nabla_s \mathrm{R}\mathrm{R}_{ij} + 3\mathrm{R}\nabla_s \mathrm{R}_{ij}\\
&\qquad +2\nabla_s|\mathrm{Ric}|^2 g_{ij} - \nabla_s \mathrm{R}^2 g_{ij}\\
&\qquad +(\nabla_i \mathrm{R}_{sp} - \nabla_p \mathrm{R}_{is})\mathrm{R}_{jp} + (\nabla_j \mathrm{R}_{sp} - \nabla_p \mathrm{R}_{js})\mathrm{R}_{ip}\\
&= \quad \nabla_s \Delta \mathrm{R}_{ij} - 5\nabla_s \mathrm{R}_{ip}\mathrm{R}_{jp} - 5\mathrm{R}_{ip}\nabla_s \mathrm{R}_{jp} + 3\nabla_s \mathrm{R}\mathrm{R}_{ij} + 3\mathrm{R}\nabla_s \mathrm{R}_{ij}\\
&\qquad +2\nabla_s|\mathrm{Ric}|^2 g_{ij} - \nabla_s \mathrm{R}^2 g_{ij} + \mathrm{C}_{spi}\mathrm{R}_{jp} + \mathrm{C}_{spj}\mathrm{R}_{ip}\\
&\qquad +\mathrm{R}_{jp}[\nabla_i \mathrm{R} g_{sp} - \nabla_p \mathrm{R} g_{is}]/4 + \mathrm{R}_{ip}[\nabla_j \mathrm{R} g_{sp} - \nabla_p \mathrm{R} g_{js}]/4\,,
\end{aligned}
$$

where in the last passage we substituted the expression of the Cotton tensor.

We then compute,

$$
\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk} &= \frac{\partial}{\partial t}\nabla_k\mathrm{R}_{ij} - \frac{\partial}{\partial t}\nabla_j\mathrm{R}_{ik} - \frac{\partial}{\partial t}\big(\nabla_k\mathrm{R}g_{ij} - \nabla_j\mathrm{R}g_{ik}\big)/4\\
&= \nabla_k\Delta\mathrm{R}_{ij} - 5\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp} - 5\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp} + 3\nabla_k\mathrm{R}\mathrm{R}_{ij} + 3\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&\quad +2\nabla_k|\mathrm{Ric}|^2 g_{ij} - \nabla_k\mathrm{R}^2 g_{ij} + \mathrm{C}_{kpi}\mathrm{R}_{jp} + \mathrm{C}_{kpj}\mathrm{R}_{ip}\\
&\quad +\mathrm{R}_{jp}[\nabla_i\mathrm{R}g_{kp} - \nabla_p\mathrm{R}g_{ik}]/4 + \mathrm{R}_{ip}[\nabla_j\mathrm{R}g_{kp} - \nabla_p\mathrm{R}g_{jk}]/4\\
&\quad -\nabla_j\Delta\mathrm{R}_{ik} + 5\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp} + 5\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp} - 3\nabla_j\mathrm{R}\mathrm{R}_{ik} - 3\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&\quad -2\nabla_j|\mathrm{Ric}|^2 g_{ik} + \nabla_j\mathrm{R}^2 g_{ik} - \mathrm{C}_{jpi}\mathrm{R}_{kp} - \mathrm{C}_{jpk}\mathrm{R}_{ip}\\
&\quad -\mathrm{R}_{kp}[\nabla_i\mathrm{R}g_{jp} - \nabla_p\mathrm{R}g_{ij}]/4 - \mathrm{R}_{ip}[\nabla_k\mathrm{R}g_{jp} - \nabla_p\mathrm{R}g_{kj}]/4\\
&\quad +\big(\mathrm{R}_{ij}\nabla_k\mathrm{R} - \mathrm{R}_{ik}\nabla_j\mathrm{R}\big)/2\\
&\quad -\big(\nabla_k\Delta\mathrm{R} + 2\nabla_k|\mathrm{Ric}|^2\big)g_{ij}/4 + \big(\nabla_j\Delta\mathrm{R} + 2\nabla_j|\mathrm{Ric}|^2\big)g_{ik}/4\\
&= \nabla_k\Delta\mathrm{R}_{ij} - 5\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp} - 5\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp} + 3\nabla_k\mathrm{R}\mathrm{R}_{ij} + 3\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&\quad +3\nabla_k|\mathrm{Ric}|^2 g_{ij}/2 - \nabla_k\mathrm{R}^2 g_{ij} + \mathrm{C}_{kpi}\mathrm{R}_{jp} + \mathrm{C}_{kpj}\mathrm{R}_{ip}\\
&\quad +\mathrm{R}_{jk}\nabla_i\mathrm{R}/4 - \mathrm{R}_{jp}\nabla_p\mathrm{R}g_{ik}/4 + \mathrm{R}_{ik}\nabla_j\mathrm{R}/4 - \mathrm{R}_{ip}\nabla_p\mathrm{R}g_{jk}/4\\
&\quad -\nabla_j\Delta\mathrm{R}_{ik} + 5\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp} + 5\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp} - 3\nabla_j\mathrm{R}\mathrm{R}_{ik} - 3\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&\quad -3\nabla_j|\mathrm{Ric}|^2 g_{ik}/2 + \nabla_j\mathrm{R}^2 g_{ik} - \mathrm{C}_{jpi}\mathrm{R}_{kp} - \mathrm{C}_{jpk}\mathrm{R}_{ip}\\
&\quad -\mathrm{R}_{kj}\nabla_i\mathrm{R}/4 + \mathrm{R}_{kp}\nabla_p\mathrm{R}g_{ij}/4 - \mathrm{R}_{ij}\nabla_k\mathrm{R}/4 + \mathrm{R}_{ip}\nabla_p\mathrm{R}g_{kj}/4\\
&\quad +\big(\mathrm{R}_{ij}\nabla_k\mathrm{R} - \mathrm{R}_{ik}\nabla_j\mathrm{R}\big)/2\\
&\quad -\nabla_k\Delta\mathrm{R}g_{ij}/4 + \nabla_j\Delta\mathrm{R}g_{ik}/4\\
&= \nabla_k\Delta\mathrm{R}_{ij} - 5\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp} - 5\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp} + 13\nabla_k\mathrm{R}\mathrm{R}_{ij}/4 + 3\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&\quad +3\nabla_k|\mathrm{Ric}|^2 g_{ij}/2 - \nabla_k\mathrm{R}^2 g_{ij} + \mathrm{C}_{kpi}\mathrm{R}_{jp} + \mathrm{C}_{kpj}\mathrm{R}_{ip}\\
&\quad -\mathrm{R}_{jp}\nabla_p\mathrm{R}g_{ik}/4\\
&\quad -\nabla_j\Delta\mathrm{R}_{ik} + 5\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp} + 5\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp} - 13\nabla_j\mathrm{R}\mathrm{R}_{ik}/4 - 3\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&\quad -3\nabla_j|\mathrm{Ric}|^2 g_{ik}/2 + \nabla_j\mathrm{R}^2 g_{ik} - \mathrm{C}_{jpi}\mathrm{R}_{kp} - \mathrm{C}_{jpk}\mathrm{R}_{ip}\\
&\quad +\mathrm{R}_{kp}\nabla_p\mathrm{R}g_{ij}/4\\
&\quad -\nabla_k\Delta\mathrm{R}g_{ij}/4 + \nabla_j\Delta\mathrm{R}g_{ik}/4
\end{aligned}
$$

and

$$
\Delta\mathrm{C}_{ijk} = \Delta\nabla_k\mathrm{R}_{ij} - \Delta\nabla_j\mathrm{R}_{ik} - \Delta\nabla_k\mathrm{R}g_{ij}/4 + \Delta\nabla_j\mathrm{R}g_{ik}/4\,,
$$

hence,

$$
\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk} - \Delta\mathrm{C}_{ijk} &= \nabla_k\Delta\mathrm{R}_{ij} - \nabla_j\Delta\mathrm{R}_{ik} - \Delta\nabla_k\mathrm{R}_{ij} + \Delta\nabla_j\mathrm{R}_{ik}\\
&\quad -\nabla_k\Delta\mathrm{R}g_{ij}/4 + \nabla_j\Delta\mathrm{R}g_{ik}/4 + \Delta\nabla_k\mathrm{R}g_{ij}/4 - \Delta\nabla_j\mathrm{R}g_{ik}/4\\
&\quad -5\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp} - 5\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp} + 13\nabla_k\mathrm{R}\mathrm{R}_{ij}/4 + 3\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&\quad +3\nabla_k|\mathrm{Ric}|^2 g_{ij}/2 - \nabla_k\mathrm{R}^2 g_{ij} + \mathrm{C}_{kpi}\mathrm{R}_{jp} + \mathrm{C}_{kpj}\mathrm{R}_{ip}\\
&\quad -\mathrm{R}_{jp}\nabla_p\mathrm{R}g_{ik}/4\\
&\quad +5\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp} + 5\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp} - 13\nabla_j\mathrm{R}\mathrm{R}_{ik}/4 - 3\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&\quad -3\nabla_j|\mathrm{Ric}|^2 g_{ik}/2 + \nabla_j\mathrm{R}^2 g_{ik} - \mathrm{C}_{jpi}\mathrm{R}_{kp} - \mathrm{C}_{jpk}\mathrm{R}_{ip}\\
&\quad +\mathrm{R}_{kp}\nabla_p\mathrm{R}g_{ij}/4
\end{aligned}
$$

Now to proceed, we need the following commutation rules for the derivatives of the Ricci tensor and of the scalar curvature, where we will employ the special form of the Riemann tensor in dimension three given by formula (2.1),

$$\begin{aligned}
\nabla_k\Delta \mathrm{R}_{ij}-\Delta\nabla_k\mathrm{R}_{ij} &= \nabla^3_{kll}\mathrm{R}_{ij}-\nabla^3_{lkl}\mathrm{R}_{ij}+\nabla^3_{lkl}\mathrm{R}_{ij}-\nabla^3_{llk}\mathrm{R}_{ij}\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\mathrm{R}_{klip}\nabla_l\mathrm{R}_{jp}+\mathrm{R}_{kljp}\nabla_l\mathrm{R}_{ip}\\
&\quad+\nabla^3_{lkl}\mathrm{R}_{ij}-\nabla^3_{llk}\mathrm{R}_{ij}\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\mathrm{R}_{ik}\nabla_j\mathrm{R}/2+\mathrm{R}_{jk}\nabla_i\mathrm{R}/2\\
&\quad-\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}-\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}+\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}+\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk}\\
&\quad-\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}-\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik}-\mathrm{R}\nabla_j\mathrm{R}g_{ik}/4-\mathrm{R}\nabla_i\mathrm{R}g_{jk}/4\\
&\quad+\mathrm{R}\nabla_i\mathrm{R}_{jk}/2+\mathrm{R}\nabla_j\mathrm{R}_{ik}/2\\
&\quad+\nabla_l\big(\mathrm{R}_{klip}\mathrm{R}_{pj}+\mathrm{R}_{kljp}\mathrm{R}_{pi}\big)\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\mathrm{R}_{ik}\nabla_j\mathrm{R}/2+\mathrm{R}_{jk}\nabla_i\mathrm{R}/2\\
&\quad-\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}-\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}+\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}+\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk}\\
&\quad-\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}-\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik}-\mathrm{R}\nabla_j\mathrm{R}g_{ik}/4-\mathrm{R}\nabla_i\mathrm{R}g_{jk}/4\\
&\quad+\mathrm{R}\nabla_i\mathrm{R}_{jk}/2+\mathrm{R}\nabla_j\mathrm{R}_{ik}/2\\
&\quad+\nabla_l\big(\mathrm{R}_{ik}\mathrm{R}_{lj}-\mathrm{R}_{il}\mathrm{R}_{kj}+\mathrm{R}_{pl}\mathrm{R}_{pj}g_{ik}-\mathrm{R}_{pk}\mathrm{R}_{pj}g_{il}-g_{ik}\mathrm{R}\mathrm{R}_{lj}/2+g_{il}\mathrm{R}\mathrm{R}_{jk}/2\\
&\quad+\mathrm{R}_{jk}\mathrm{R}_{li}-\mathrm{R}_{jl}\mathrm{R}_{ki}+\mathrm{R}_{pl}\mathrm{R}_{pi}g_{jk}-\mathrm{R}_{pk}\mathrm{R}_{pi}g_{jl}-g_{jk}\mathrm{R}\mathrm{R}_{li}/2+g_{jl}\mathrm{R}\mathrm{R}_{ik}/2\big)\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\mathrm{R}_{ik}\nabla_j\mathrm{R}/2+\mathrm{R}_{jk}\nabla_i\mathrm{R}/2\\
&\quad-\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}-\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}+\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}+\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk}\\
&\quad-\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}-\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik}-\mathrm{R}\nabla_j\mathrm{R}g_{ik}/4-\mathrm{R}\nabla_i\mathrm{R}g_{jk}/4\\
&\quad+\mathrm{R}\nabla_i\mathrm{R}_{jk}/2+\mathrm{R}\nabla_j\mathrm{R}_{ik}/2\\
&\quad-\nabla_i\mathrm{R}_{pk}\mathrm{R}_{pj}+\nabla_i\mathrm{R}\mathrm{R}_{jk}/2+g_{ik}\mathrm{R}_{pl}\nabla_l\mathrm{R}_{pj}\\
&\quad-\mathrm{R}_{pk}\nabla_i\mathrm{R}_{pj}-g_{ik}\mathrm{R}\nabla_j\mathrm{R}/4+\mathrm{R}\nabla_i\mathrm{R}_{jk}/2\\
&\quad-\nabla_j\mathrm{R}_{pk}\mathrm{R}_{pi}+\nabla_j\mathrm{R}\mathrm{R}_{ik}/2+g_{jk}\mathrm{R}_{pl}\nabla_l\mathrm{R}_{pi}\\
&\quad-\mathrm{R}_{pk}\nabla_j\mathrm{R}_{pi}-g_{jk}\mathrm{R}\nabla_i\mathrm{R}/4+\mathrm{R}\nabla_j\mathrm{R}_{ik}/2\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\mathrm{R}_{ik}\nabla_j\mathrm{R}+\mathrm{R}_{jk}\nabla_i\mathrm{R}\\
&\quad-2\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}-2\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}+2\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}+2\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk}\\
&\quad-\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}-\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik}-\mathrm{R}_{pj}\nabla_i\mathrm{R}_{pk}-\mathrm{R}_{pi}\nabla_j\mathrm{R}_{pk}\\
&\quad-\mathrm{R}\nabla_j\mathrm{R}g_{ik}/2-\mathrm{R}\nabla_i\mathrm{R}g_{jk}/2+\mathrm{R}\nabla_i\mathrm{R}_{jk}+\mathrm{R}\nabla_j\mathrm{R}_{ik}
\end{aligned}$$

and

$$\nabla_k\Delta\mathrm{R}-\Delta\nabla_k\mathrm{R}=\mathrm{R}_{kllp}\nabla_p\mathrm{R}=-\mathrm{R}_{kp}\nabla_p\mathrm{R}\,.$$

Then, getting back to the main computation, we obtain

$$
\begin{aligned}
\frac{\partial}{\partial t}C_{ijk} - \Delta C_{ijk} \;=\;& -R_{kp}\nabla_p R_{ij} + R_{ik}\nabla_j R + R_{jk}\nabla_i R\\
& -2R_{kp}\nabla_i R_{jp} - 2R_{kp}\nabla_j R_{ip} + 2R_{lp}\nabla_l R_{jp}g_{ik} + 2R_{lp}\nabla_l R_{ip}g_{jk}\\
& -R_{li}\nabla_l R_{jk} - R_{lj}\nabla_l R_{ik} - R_{pj}\nabla_i R_{pk} - R_{pi}\nabla_j R_{pk}\\
& -R\nabla_j R g_{ik}/2 - R\nabla_i R g_{jk}/2 + R\nabla_i R_{jk} + R\nabla_j R_{ik}\\
& +R_{jp}\nabla_p R_{ik} - R_{ij}\nabla_k R - R_{kj}\nabla_i R\\
& +2R_{jp}\nabla_i R_{kp} + 2R_{jp}\nabla_k R_{ip} - 2R_{lp}\nabla_l R_{kp}g_{ij} - 2R_{lp}\nabla_l R_{ip}g_{kj}\\
& +R_{li}\nabla_l R_{kj} + R_{lk}\nabla_l R_{ij} + R_{pk}\nabla_i R_{pj} + R_{pi}\nabla_k R_{pj}\\
& +R\nabla_k R g_{ij}/2 + R\nabla_i R g_{kj}/2 - R\nabla_i R_{kj} - R\nabla_k R_{ij}\\
& +R_{kp}\nabla_p R g_{ij}/4 - R_{jp}\nabla_p R g_{ik}/4\\
& -5\nabla_k R_{ip}R_{jp} - 5R_{ip}\nabla_k R_{jp} + 13\nabla_k R R_{ij}/4 + 3R\nabla_k R_{ij}\\
& +3\nabla_k|\mathrm{Ric}|^2 g_{ij}/2 - \nabla_k R^2 g_{ij} + C_{kpi}R_{jp} + C_{kpj}R_{ip}\\
& -R_{jp}\nabla_p R g_{ik}/4\\
& +5\nabla_j R_{ip}R_{kp} + 5R_{ip}\nabla_j R_{kp} - 13\nabla_j R R_{ik}/4 - 3R\nabla_j R_{ik}\\
& -3\nabla_j|\mathrm{Ric}|^2 g_{ik}/2 + \nabla_j R^2 g_{ik} - C_{jpi}R_{kp} - C_{jpk}R_{ip}\\
& +R_{kp}\nabla_p R g_{ij}/4\\
=\;& C_{kpi}R_{jp} + C_{kpj}R_{ip} - C_{jpi}R_{kp} - C_{jpk}R_{ip}\\
& +[2R_{lp}\nabla_l R_{jp} + 3R\nabla_j R/2 - R_{jp}\nabla_p R/2 - 3\nabla_j|\mathrm{Ric}|^2/2]g_{ik}\\
& +[-2R_{lp}\nabla_l R_{kp} - 3R\nabla_k R/2 + R_{kp}\nabla_p R/2 + 3\nabla_k|\mathrm{Ric}|^2/2]g_{ij}\\
& -R_{kp}\nabla_i R_{jp} + R_{jp}\nabla_i R_{kp}\\
& -3\nabla_k R_{ip}R_{jp} - 4R_{ip}\nabla_k R_{jp} + 9\nabla_k R R_{ij}/4 + 2R\nabla_k R_{ij}\\
& +3\nabla_j R_{ip}R_{kp} + 4R_{ip}\nabla_j R_{kp} - 9\nabla_j R R_{ik}/4 - 2R\nabla_j R_{ik}
\end{aligned}
$$

Now, by means of the very definition of the Cotton tensor in dimension three (2.2) and the identities (1.4), we substitute

$$
\begin{aligned}
C_{kpj} - C_{jpk} = -\,C_{kjp} - C_{jpk} &= C_{pkj}\\
\nabla_l R_{jp} &= \nabla_j R_{lp} + C_{pjl} + \frac{1}{4}\big(\nabla_l R g_{pj} - \nabla_j R g_{pl}\big)\\
\nabla_l R_{kp} &= \nabla_k R_{lp} + C_{pkl} + \frac{1}{4}\big(\nabla_l R g_{pk} - \nabla_k R g_{pl}\big)\\
\nabla_i R_{jp} &= \nabla_j R_{ip} + C_{pji} + \frac{1}{4}\big(\nabla_i R g_{jp} - \nabla_j R g_{ip}\big)\\
\nabla_i R_{kp} &= \nabla_k R_{ip} + C_{pki} + \frac{1}{4}\big(\nabla_i R g_{kp} - \nabla_k R g_{ip}\big)
\end{aligned}
$$

in the last expression above, getting

$$
\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk}-\Delta\mathrm{C}_{ijk} \;=\;& \mathrm{R}_{jp}\mathrm{C}_{kpi}-\mathrm{R}_{kp}\mathrm{C}_{jpi}+\mathrm{R}_{ip}\mathrm{C}_{pkj}\\
&+\Big[2\mathrm{R}_{lp}\big(\nabla_j\mathrm{R}_{lp}+\mathrm{C}_{pjl}+\nabla_l\mathrm{R}g_{pj}/4-\nabla_j\mathrm{R}g_{pl}/4\big)\\
&\qquad+3\mathrm{R}\nabla_j\mathrm{R}/2-\mathrm{R}_{jp}\nabla_p\mathrm{R}/2-3\nabla_j|\mathrm{Ric}|^2/2\Big]g_{ik}\\
&+\Big[-2\mathrm{R}_{lp}\big(\nabla_k\mathrm{R}_{lp}+\mathrm{C}_{pkl}+\nabla_l\mathrm{R}g_{pk}/4-\nabla_k\mathrm{R}g_{pl}/4\big)\\
&\qquad-3\mathrm{R}\nabla_k\mathrm{R}/2+\mathrm{R}_{kp}\nabla_p\mathrm{R}/2+3\nabla_k|\mathrm{Ric}|^2/2\Big]g_{ij}\\
&-\mathrm{R}_{kp}\big(\nabla_j\mathrm{R}_{ip}+\mathrm{C}_{pji}+\nabla_i\mathrm{R}g_{jp}/4-\nabla_j\mathrm{R}g_{ip}/4\big)\\
&+\mathrm{R}_{jp}\big(\nabla_k\mathrm{R}_{ip}+\mathrm{C}_{pki}+\nabla_i\mathrm{R}g_{kp}/4-\nabla_k\mathrm{R}g_{ip}/4\big)\\
&-3\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp}-4\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp}+9\nabla_k\mathrm{R}\mathrm{R}_{ij}/4+2\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&+3\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp}+4\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp}-9\nabla_j\mathrm{R}\mathrm{R}_{ik}/4-2\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
\;=\;& \mathrm{R}_{jp}\big(\mathrm{C}_{kpi}+\mathrm{C}_{pki}\big)-\mathrm{R}_{kp}\big(\mathrm{C}_{jpi}+\mathrm{C}_{pji}\big)+\mathrm{R}_{ip}\mathrm{C}_{pkj}\\
&+2\mathrm{R}_{lp}\mathrm{C}_{pjl}g_{ik}-2\mathrm{R}_{lp}\mathrm{C}_{pkl}g_{ij}\\
&+\big[\mathrm{R}\nabla_j\mathrm{R}-\nabla_j|\mathrm{Ric}|^2/2\big]g_{ik}-\big[\mathrm{R}\nabla_k\mathrm{R}-\nabla_k|\mathrm{Ric}|^2/2\big]g_{ij}\\
&-2\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp}-4\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp}+2\nabla_k\mathrm{R}\mathrm{R}_{ij}+2\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&+2\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp}+4\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp}-2\nabla_j\mathrm{R}\mathrm{R}_{ik}-2\mathrm{R}\nabla_j\mathrm{R}_{ik}\,.
\end{aligned}
$$

then, we substitute again

$$
\begin{aligned}
\nabla_k\mathrm{R}_{jp} &= \nabla_p\mathrm{R}_{kj}+\mathrm{C}_{jpk}+\frac{1}{4}\big(\nabla_k\mathrm{R}g_{jp}-\nabla_p\mathrm{R}g_{jk}\big)\\
\nabla_j\mathrm{R}_{kp} &= \nabla_p\mathrm{R}_{jk}+\mathrm{C}_{kpj}+\frac{1}{4}\big(\nabla_j\mathrm{R}g_{kp}-\nabla_p\mathrm{R}g_{kj}\big)\\
\nabla_k\mathrm{R}_{ij} &= \nabla_i\mathrm{R}_{kj}+\mathrm{C}_{jik}+\frac{1}{4}\big(\nabla_k\mathrm{R}g_{ij}-\nabla_i\mathrm{R}g_{jk}\big)\\
\nabla_j\mathrm{R}_{ik} &= \nabla_i\mathrm{R}_{jk}+\mathrm{C}_{kij}+\frac{1}{4}\big(\nabla_j\mathrm{R}g_{ik}-\nabla_i\mathrm{R}g_{kj}\big)\,,
\end{aligned}
$$

finally obtaining

$$\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk}-\Delta\mathrm{C}_{ijk} =\ & \mathrm{R}_{jp}\big(\mathrm{C}_{kpi}+\mathrm{C}_{pki}\big)-\mathrm{R}_{kp}\big(\mathrm{C}_{jpi}+\mathrm{C}_{pji}\big)+\mathrm{R}_{ip}\mathrm{C}_{pkj}\\
&+2\mathrm{R}_{lp}\mathrm{C}_{pjl}g_{ik}-2\mathrm{R}_{lp}\mathrm{C}_{pkl}g_{ij}\\
&+\big[\mathrm{R}\nabla_j\mathrm{R}-\nabla_j|\mathrm{Ric}|^2/2\big]g_{ik}-\big[\mathrm{R}\nabla_k\mathrm{R}-\nabla_k|\mathrm{Ric}|^2/2\big]g_{ij}\\
&-2\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp}-4\mathrm{R}_{ip}\big(\nabla_p\mathrm{R}_{kj}+\mathrm{C}_{jpk}+\nabla_k\mathrm{R}g_{jp}/4-\nabla_p\mathrm{R}g_{jk}/4\big)\\
&+2\nabla_k\mathrm{R}\mathrm{R}_{ij}+2\mathrm{R}\big(\nabla_i\mathrm{R}_{kj}+\mathrm{C}_{jik}+\nabla_k\mathrm{R}g_{ij}/4-\nabla_i\mathrm{R}g_{jk}/4\big)\\
&+2\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp}+4\mathrm{R}_{ip}\big(\nabla_p\mathrm{R}_{jk}+\mathrm{C}_{kpj}+\nabla_j\mathrm{R}g_{kp}/4-\nabla_p\mathrm{R}g_{kj}/4\big)\\
&-2\nabla_j\mathrm{R}\mathrm{R}_{ik}-2\mathrm{R}\big(\nabla_i\mathrm{R}_{jk}+\mathrm{C}_{kij}+\nabla_j\mathrm{R}g_{ik}/4-\nabla_i\mathrm{R}g_{kj}/4\big)\\
=\ & \mathrm{R}_{jp}\big(\mathrm{C}_{kpi}+\mathrm{C}_{pki}\big)-\mathrm{R}_{kp}\big(\mathrm{C}_{jpi}+\mathrm{C}_{pji}\big)+\mathrm{R}_{ip}\mathrm{C}_{pkj}\\
&+4\mathrm{R}_{ip}\big(\mathrm{C}_{kpj}-\mathrm{C}_{jpk}\big)+2\mathrm{R}\big(\mathrm{C}_{jik}-\mathrm{C}_{kij}\big)\\
&+2\mathrm{R}_{lp}\mathrm{C}_{pjl}g_{ik}-2\mathrm{R}_{lp}\mathrm{C}_{pkl}g_{ij}\\
&+\big[\mathrm{R}\nabla_j\mathrm{R}/2-\nabla_j|\mathrm{Ric}|^2/2\big]g_{ik}-\big[\mathrm{R}\nabla_k\mathrm{R}/2-\nabla_k|\mathrm{Ric}|^2/2\big]g_{ij}\\
&-2\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp}+2\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp}\\
&+\nabla_k\mathrm{R}\mathrm{R}_{ij}-\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
=\ & \mathrm{R}_{jp}\big(\mathrm{C}_{kpi}+\mathrm{C}_{pki}\big)-\mathrm{R}_{kp}\big(\mathrm{C}_{jpi}+\mathrm{C}_{pji}\big)+5\mathrm{R}_{ip}\mathrm{C}_{pkj}\\
&+2\mathrm{R}\mathrm{C}_{ijk}+2\mathrm{R}_{lp}\mathrm{C}_{pjl}g_{ik}-2\mathrm{R}_{lp}\mathrm{C}_{pkl}g_{ij}\\
&+\big[\mathrm{R}\nabla_j\mathrm{R}/2-\nabla_j|\mathrm{Ric}|^2/2\big]g_{ik}-\big[\mathrm{R}\nabla_k\mathrm{R}/2-\nabla_k|\mathrm{Ric}|^2/2\big]g_{ij}\\
&+2\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp}-2\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp}\\
&+\nabla_k\mathrm{R}\mathrm{R}_{ij}-\nabla_j\mathrm{R}\mathrm{R}_{ik}\,,
\end{aligned}$$

where in the last passage we used again the identities (1.4).
Hence, we can resume this long computation in the following proposition, getting back to a generic coordinate basis.

**Proposition 2.1.** *During the Ricci flow of a 3–dimensional Riemannian manifold $(M^3,g(t))$, the Cotton tensor satisfies the following evolution equation*

$$\begin{aligned}
(\partial_t-\Delta)\mathrm{C}_{ijk} =\ & g^{pq}\mathrm{R}_{pj}(\mathrm{C}_{kqi}+\mathrm{C}_{qki})+5g^{pq}\mathrm{R}_{ip}\mathrm{C}_{qkj}+g^{pq}\mathrm{R}_{pk}(\mathrm{C}_{jiq}+\mathrm{C}_{qij})\\
&+2\mathrm{R}\mathrm{C}_{ijk}+2\mathrm{R}^{ql}\mathrm{C}_{qjl}g_{ik}-2\mathrm{R}^{ql}\mathrm{C}_{qkl}g_{ij}\\
&+\frac{1}{2}\nabla_k|\mathrm{Ric}|^2g_{ij}-\frac{1}{2}\nabla_j|\mathrm{Ric}|^2g_{ik}+\frac{\mathrm{R}}{2}\nabla_j\mathrm{R}g_{ik}-\frac{\mathrm{R}}{2}\nabla_k\mathrm{R}g_{ij}\\
&+2g^{pq}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{qi}-2g^{pq}\mathrm{R}_{pj}\nabla_k\mathrm{R}_{qi}+\mathrm{R}_{ij}\nabla_k\mathrm{R}-\mathrm{R}_{ik}\nabla_j\mathrm{R}\,.
\end{aligned}\tag{2.3}$$

*In particular if the Cotton tensor vanishes identically along the flow we obtain,*

$$\begin{aligned}
0 =\ & \nabla_k|\mathrm{Ric}|^2g_{ij}-\nabla_j|\mathrm{Ric}|^2g_{ik}+\mathrm{R}\nabla_j\mathrm{R}g_{ik}-\mathrm{R}\nabla_k\mathrm{R}g_{ij}\\
&+4g^{pq}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{qi}-4g^{pq}\mathrm{R}_{pj}\nabla_k\mathrm{R}_{qi}+2\mathrm{R}_{ij}\nabla_k\mathrm{R}-2\mathrm{R}_{ik}\nabla_j\mathrm{R}\,.
\end{aligned}\tag{2.4}$$

**Corollary 2.2.** *If the Cotton tensor vanishes identically along the Ricci flow of a 3–dimensional Riemannian manifold $(M^3,g(t))$, the following tensor*

$$|\mathrm{Ric}|^2g_{ij}-4\mathrm{R}_{pj}\mathrm{R}_{pi}+3\mathrm{R}\mathrm{R}_{ij}-\frac{7}{8}\mathrm{R}^2g_{ij}$$

*is a Codazzi tensor (see* [1, Chapter 16, Section C]*).*

*Proof.* We compute in an orthonormal basis,

$$
\begin{aligned}
4R_{pk}\nabla_j R_{pi} - 4\,R_{pj}\nabla_k R_{pi} + 2R_{ij}\nabla_k R - 2R_{ik}\nabla_j R
&= 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi}) - 4R_{pi}\nabla_j R_{pk} + 4R_{pi}\nabla_k R_{pj}\\
&\quad + 2R_{ij}\nabla_k R - 2R_{ik}\nabla_j R\\
&= 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi}) + R_{pi}(4C_{pjk} + \nabla_k R g_{pj} - \nabla_j R g_{pk})\\
&\quad + 2R_{ij}\nabla_k R - 2R_{ik}\nabla_j R\\
&= 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi}) + 3R_{ij}\nabla_k R - 3R_{ik}\nabla_j R\\
&= 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi}) + 3\nabla_k(RR_{ij}) - 3\nabla_j(RR_{ik})\\
&\quad - 3R(\nabla_k R_{ij} - \nabla_j R_{ik})\\
&= 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi}) + 3\nabla_k(RR_{ij}) - 3\nabla_j(RR_{ik})\\
&\quad - 3R(4C_{ijk} + \nabla_k R g_{ij} - \nabla_j R g_{ik})/4\\
&= 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi}) + 3\nabla_k(RR_{ij}) - 3\nabla_j(RR_{ik})\\
&\quad - \frac{3}{8}\nabla_k R^2 g_{ij} + \frac{3}{8}\nabla_j R^2 g_{ik}\,.
\end{aligned}
$$

Hence, we have, by the previous proposition,

$$
\begin{aligned}
0 &= \nabla_k|\mathrm{Ric}|^2 g_{ij} - \nabla_j|\mathrm{Ric}|^2 g_{ik} + 4\nabla_j(R_{pk}R_{pi}) - 4\nabla_k(R_{pj}R_{pi})\\
&\quad + 3\nabla_k(RR_{ij}) - 3\nabla_j(RR_{ik}) - \frac{7}{8}\nabla_k R^2 g_{ij} + \frac{7}{8}\nabla_j R^2 g_{ik}\,,
\end{aligned}
$$

which is the thesis of the corollary. □

*Remark* 2.3. All the traces of the 3–tensor on the LHS of equation (2.4) are zero.

*Remark* 2.4. From the trace–free property (1.5) of the Cotton tensor and the fact that along the Ricci flow there holds

$$
(\partial_t - \Delta)g^{ij} = 2R^{ij}\,,
$$

we conclude that the following relations have to hold

$$
\begin{aligned}
g^{ij}(\partial_t - \Delta)C_{ijk} &= -2R^{ij}C_{ijk}\,,\\
g^{ik}(\partial_t - \Delta)C_{ijk} &= -2R^{ik}C_{ijk}\,,\\
g^{jk}(\partial_t - \Delta)C_{ijk} &= -2R^{jk}C_{ijk} = 0\,.
\end{aligned}
$$

They are easily verified for formula (2.3).

**Corollary 2.5.** *During the Ricci flow of a 3–dimensional Riemannian manifold $(M^3, g(t))$, the squared norm of the Cotton tensor satisfies the following evolution equation, in an orthonormal basis,*

$$
\begin{aligned}
(\partial_t - \Delta)|C_{ijk}|^2 &= -2|\nabla C_{ijk}|^2 - 24C_{ipk}C_{iqk}R_{pq} + 28C_{ipk}C_{kqi}R_{pq} + 4R|C_{ijk}|^2\\
&\quad + 8C_{ijk}R_{pk}\nabla_j R_{pi} + 4C_{ijk}R_{ij}\nabla_k R\,.
\end{aligned}
$$

*Proof.*

$$
\begin{aligned}
(\partial_t - \Delta)|C_{ijk}|^2 &= -2|\nabla C_{ijk}|^2 + 2C^{ijk}\Big[g^{pq}R_{pj}(C_{kqi} + C_{qki}) + 5g^{pq}R_{ip}C_{qkj} + g^{pq}R_{pk}(C_{jiq} + C_{qij})\\
&\quad + 2RC_{ijk} + 2R^{ql}C_{qjl}g_{ik} - 2R^{ql}C_{qkl}g_{ij}\\
&\quad + \frac{1}{2}\nabla_k|\mathrm{Ric}|^2 g_{ij} - \frac{1}{2}\nabla_j|\mathrm{Ric}|^2 g_{ik} + \frac{R}{2}\nabla_j R g_{ik} - \frac{R}{2}\nabla_k R g_{ij}\\
&\quad + 2g^{pq}R_{pk}\nabla_j R_{qi} - 2g^{pq}R_{pj}\nabla_k R_{qi} + R_{ij}\nabla_k R - R_{ik}\nabla_j R\Big]\\
&= -2|\nabla C_{ijk}|^2 + 2C^{ijk}\Big[2g^{pq}R_{pj}(C_{kqi} + C_{qki}) + 5g^{pq}R_{ip}C_{qkj}\Big]\\
&\quad + 4R|C_{ijk}|^2 + 8g^{pq}C^{ijk}R_{pk}\nabla_j R_{qi} + 4C^{ijk}R_{ij}\nabla_k R\\
&= -2|\nabla C_{ijk}|^2 - 24C_{ipk}C_{iqk}R_{pq} + 28C_{ipk}C_{kqi}R_{pq} + 4R|C_{ijk}|^2\\
&\quad + 8C_{ijk}R_{pk}\nabla_j R_{pi} + 4C_{ijk}R_{ij}\nabla_k R\,.
\end{aligned}
$$

□

## 3. THREE–DIMENSIONAL GRADIENT RICCI SOLITONS

The structural equation of a gradient Ricci soliton $(M, g)$ is the following

$$\mathrm{R}_{ij} + \nabla_i\nabla_j f = \lambda g_{ij}\,, \tag{3.1}$$

for some $\lambda \in \mathbb{R}$.
The soliton is said to be *steady*, *shrinking* or *expanding* according to the fact that the constant $\lambda$ is zero, positive or negative, respectively.

It follows that in dimension three, for $(M^3, g)$ there holds

$$\begin{aligned}
\Delta\mathrm{R}_{ij} &= \nabla_l\mathrm{R}_{ij}\nabla_l f + 2\lambda\mathrm{R}_{ij} - 2|\mathrm{Ric}|^2 g_{ij} + \mathrm{R}^2 g_{ij} - 3\mathrm{R}\mathrm{R}_{ij} + 4\mathrm{R}_{is}\mathrm{R}_{sj} && (3.2)\\
\Delta\mathrm{R} &= \nabla_l\mathrm{R}\nabla_l f + 2\lambda\mathrm{R} - 2|\mathrm{Ric}|^2 && (3.3)\\
\nabla_i\mathrm{R} &= 2\mathrm{R}_{li}\nabla_l f && (3.4)\\
\mathrm{C}_{ijk} &= \frac{\mathrm{R}_{lk}g_{ij}}{2}\nabla_l f - \frac{\mathrm{R}_{lj}g_{ik}}{2}\nabla_l f + \mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f + \frac{\mathrm{R}g_{ik}}{2}\nabla_j f - \frac{\mathrm{R}g_{ij}}{2}\nabla_k f && (3.5)\\
&= \frac{\nabla_k\mathrm{R}}{4}g_{ij} - \frac{\nabla_j\mathrm{R}}{4}g_{ik} + \Big(\mathrm{R}_{ij} - \frac{\mathrm{R}}{2}g_{ij}\Big)\nabla_k f - \Big(\mathrm{R}_{ik} - \frac{\mathrm{R}}{2}g_{ik}\Big)\nabla_j f\,.
\end{aligned}$$

In the special case of a *steady* soliton the first two equations above simplify as follows,

$$\begin{aligned}
\Delta\mathrm{R}_{ij} &= \nabla_l\mathrm{R}_{ij}\nabla_l f - 2|\mathrm{Ric}|^2 g_{ij} + \mathrm{R}^2 g_{ij} - 3\mathrm{R}\mathrm{R}_{ij} + 4\mathrm{R}_{is}\mathrm{R}_{sj}\\
\Delta\mathrm{R} &= \nabla_l\mathrm{R}\nabla_l f - 2|\mathrm{Ric}|^2\,.
\end{aligned}$$

*Remark* 3.1. We notice that, by relation (3.5), we have

$$\begin{aligned}
\mathrm{C}_{ijk}\nabla_i f &= \frac{\nabla_k\mathrm{R}\nabla_j f}{4} - \frac{\nabla_j\mathrm{R}\nabla_k f}{4} + \mathrm{R}_{ij}\nabla_i f\nabla_k f - \frac{\mathrm{R}}{2}\nabla_j f\nabla_k f - \mathrm{R}_{ik}\nabla_i f\nabla_j f + \frac{\mathrm{R}}{2}\nabla_k f\nabla_j f\\
&= \frac{\nabla_j\mathrm{R}\nabla_k f}{4} - \frac{\nabla_k\mathrm{R}\nabla_j f}{4}\,,
\end{aligned}$$

where in the last passage we used relation (3.4).
It follows that

$$\mathrm{C}_{ijk}\nabla_i f\nabla_j f = \frac{\langle\nabla f, \nabla\mathrm{R}\rangle}{4}\nabla_k f - \frac{|\nabla f|^2}{4}\nabla_k\mathrm{R}\,.$$

Hence, if the Cotton tensor of a three–dimensional gradient Ricci soliton is identically zero, we have that at every point where $\nabla\mathrm{R}$ is not zero, $\nabla f$ and $\nabla\mathrm{R}$ are proportional.

This relation is a key step in (yet another) proof of the fact that a three–dimensional, locally conformally flat, gradient Ricci soliton is locally a warped product of a constant curvature surface on a interval of $\mathbb{R}$, leading to a full classification, first obtained by H.-D. Cao and Q. Chen [3] for the steady case and H.-D. Cao, B.-L. Chen and X.-P. Zhu [2] for the shrinking case (actually this is the last paper of a series finally classifying, in full generality, all the three-dimensional gradient Ricci solitons, even without the LCF assumption).

By a straightforward computation we have the following proposition.

**Proposition 3.2.** *Let $(M^3, g)$ be a 3–dimensional gradient Ricci soliton. Then,*

$$\begin{aligned}
\Delta|\mathrm{C}_{ijk}|^2 = {}& \nabla_l|\mathrm{C}_{ijk}|^2\nabla_l f + 2|\nabla\mathrm{C}_{ijk}|^2 + 6\lambda|\mathrm{C}_{ijk}|^2 - 2\mathrm{R}|\mathrm{C}_{ijk}|^2\\
&-6\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k\mathrm{R} + 8\mathrm{C}_{jsk}\mathrm{C}_{jik}\mathrm{R}_{si} - 16\mathrm{C}_{jsk}\mathrm{C}_{kij}\mathrm{R}_{si} - 8\mathrm{C}_{ijk}\mathrm{R}_{lk}\nabla_j\mathrm{R}_{il}\,.
\end{aligned} \tag{3.6}$$

*Hence, by Corollary 2.5, we obtain that during the Ricci flow of a 3–dimensional gradient Ricci soliton*

$$\begin{aligned}
\partial_t|\mathrm{C}_{ijk}|^2 = {}& \nabla_l|\mathrm{C}_{ijk}|^2\nabla_l f + 6\lambda|\mathrm{C}_{ijk}|^2 + 2\mathrm{R}|\mathrm{C}_{ijk}|^2\\
&-2\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k\mathrm{R} - 16\mathrm{C}_{jsk}\mathrm{C}_{jik}\mathrm{R}_{si} + 12\mathrm{C}_{jsk}\mathrm{C}_{kij}\mathrm{R}_{si}\,.
\end{aligned} \tag{3.7}$$

*Proof.* First observe that

$$\Delta|\mathrm{C}_{ijk}|^2 = 2\mathrm{C}_{ijk}\Delta\mathrm{C}_{ijk} + 2|\nabla\mathrm{C}_{ijk}|^2.$$

Using relations (3.5), (3.2) and, repeatedly, the trace-free property (1.5) of the Cotton tensor, we have that

$$\begin{aligned}
\mathrm{C}_{ijk}\Delta\mathrm{C}_{ijk} &= \Delta(\mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f)\mathrm{C}_{ijk}\\
&= (\Delta\mathrm{R}_{ij}\nabla_k f + \mathrm{R}_{ij}\Delta\nabla_k f + 2\nabla_l\mathrm{R}_{ij}\nabla_l\nabla_k f)\mathrm{C}_{ijk}\\
&\quad -(\Delta\mathrm{R}_{ik}\nabla_j f + \mathrm{R}_{ik}\Delta\nabla_j f + 2\nabla_l\mathrm{R}_{ik}\nabla_l\nabla_j f)\mathrm{C}_{ijk}\\
&= (\nabla_s\mathrm{R}_{ij}\nabla_s f + 2\lambda\mathrm{R}_{ij} - 3\mathrm{R}\mathrm{R}_{ij} + 4\mathrm{R}_{is}\mathrm{R}_{sj})\nabla_k f\mathrm{C}_{ijk}\\
&\quad +\mathrm{R}_{ij}\Delta\nabla_k f\mathrm{C}_{ijk} + 2\nabla_l\mathrm{R}_{ij}\nabla_l\nabla_k f\mathrm{C}_{ijk}\\
&\quad -(\nabla_s\mathrm{R}_{ik}\nabla_s f + 2\lambda\mathrm{R}_{ik} - 3\mathrm{R}\mathrm{R}_{ik} + 4\mathrm{R}_{is}\mathrm{R}_{sk})\nabla_j f\mathrm{C}_{ijk}\\
&\quad -\mathrm{R}_{ik}\Delta\nabla_j f\mathrm{C}_{ijk} - 2\nabla_l\mathrm{R}_{ik}\nabla_l\nabla_j f\mathrm{C}_{ijk}\\
&= (\nabla_s\mathrm{R}_{ij}\nabla_k f - \nabla_s\mathrm{R}_{ik}\nabla_j f)\nabla_s f\mathrm{C}_{ijk}\\
&\quad +2\lambda(\mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f)\mathrm{C}_{ijk} - 3\mathrm{R}(\mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f)\mathrm{C}_{ijk}\\
&\quad +4\mathrm{R}_{is}(\mathrm{R}_{sj}\nabla_k f - \mathrm{R}_{sk}\nabla_j f)\mathrm{C}_{ijk}\\
&\quad +(\mathrm{R}_{ij}\nabla_l\nabla_l\nabla_k f - \mathrm{R}_{ik}\nabla_l\nabla_l\nabla_j f)\mathrm{C}_{ijk}\\
&\quad +2(\nabla_l\mathrm{R}_{ij}\nabla_l\nabla_k f - \nabla_l\mathrm{R}_{ik}\nabla_l\nabla_j f)\mathrm{C}_{ijk}\\
&= (\nabla_s\mathrm{R}_{ij}\nabla_k f - \nabla_s\mathrm{R}_{ik}\nabla_j f)\nabla_s f\mathrm{C}_{ijk}\\
&\quad +(2\lambda - 3\mathrm{R})|\mathrm{C}_{ijk}|^2\\
&\quad +4\mathrm{R}_{is}(\mathrm{R}_{sj}\nabla_k f - \mathrm{R}_{sk}\nabla_j f)\mathrm{C}_{ijk}\\
&\quad +(\mathrm{R}_{ij}\nabla_l\nabla_l\nabla_k f - \mathrm{R}_{ik}\nabla_l\nabla_l\nabla_j f)\mathrm{C}_{ijk}\\
&\quad +2(\nabla_l\mathrm{R}_{ij}\nabla_l\nabla_k f - \nabla_l\mathrm{R}_{ik}\nabla_l\nabla_j f)\mathrm{C}_{ijk}\,,
\end{aligned}$$

where we used the identity

$$(\mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f)\mathrm{C}_{ijk} = |\mathrm{C}_{ijk}|^2$$

which follows easily by equation (3.5).
Using now equations (3.1), (3.5), (1.5), (1.4), and (3.4), we compute

$$\begin{aligned}
(\nabla_s \mathrm{R}_{ij}\nabla_k f - \nabla_s \mathrm{R}_{ik}\nabla_j f)\nabla_s f \mathrm{C}_{ijk} &= (\nabla_s(\mathrm{R}_{ij}\nabla_k f) - \mathrm{R}_{ij}\nabla_s\nabla_k f)\nabla_s f \mathrm{C}_{ijk} \\
&\quad -(\nabla_s(\mathrm{R}_{ik}\nabla_j f) - \mathrm{R}_{ik}\nabla_s\nabla_j f)\nabla_s f \mathrm{C}_{ijk} \\
&= (\nabla_s(\mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f) + \mathrm{R}_{ij}(\mathrm{R}_{sk} - \lambda g_{sk}))\nabla_s f \mathrm{C}_{ijk} \\
&\quad -(\mathrm{R}_{ik}(\mathrm{R}_{sj} - \lambda g_{sj}))\nabla_s f \mathrm{C}_{ijk} \\
&= \nabla_s \mathrm{C}_{ijk}\mathrm{C}_{ijk}\nabla_s f + \mathrm{R}_{ij}\mathrm{R}_{sk}\nabla_s f \mathrm{C}_{ijk} - \mathrm{R}_{ik}\mathrm{R}_{sj}\nabla_s f \mathrm{C}_{ijk} \\
&\quad -\lambda(\mathrm{R}_{ij}\nabla_k f - \mathrm{R}_{ik}\nabla_j f)\mathrm{C}_{ijk} \\
&= \frac{1}{2}\nabla_s|\mathrm{C}_{ijk}|^2\nabla_s f + \frac{1}{2}\mathrm{R}_{ij}\nabla_k \mathrm{R}\mathrm{C}_{ijk} - \frac{1}{2}\mathrm{R}_{ik}\nabla_j \mathrm{R}\mathrm{C}_{ijk} - \lambda|\mathrm{C}_{ijk}|^2
\end{aligned}$$

$$\begin{aligned}
4\mathrm{R}_{is}(\mathrm{R}_{sj}\nabla_k f - \mathrm{R}_{sk}\nabla_j f)\mathrm{C}_{ijk} &= 4\mathrm{R}_{is}(\mathrm{C}_{sjk} - \frac{1}{4}\nabla_k \mathrm{R} g_{sj} + \frac{1}{4}\nabla_j \mathrm{R} g_{sk} + \frac{\mathrm{R}}{2}\nabla_k f g_{sj} - \frac{\mathrm{R}}{2}\nabla_j f g_{sk})\mathrm{C}_{ijk} \\
&= 4\mathrm{R}_{is}(-\mathrm{C}_{jks} - \mathrm{C}_{ksj})(-\mathrm{C}_{jki} - \mathrm{C}_{kij}) - \mathrm{R}_{ij}\nabla_k \mathrm{R}\mathrm{C}_{ijk} \\
&\quad +\mathrm{R}_{ik}\nabla_j \mathrm{R}\mathrm{C}_{ijk} + 2\mathrm{R}\mathrm{R}_{ij}\nabla_k f \mathrm{C}_{ijk} - 2\mathrm{R}\mathrm{R}_{ik}\nabla_j f \mathrm{C}_{ijk} \\
&= 8\mathrm{R}_{is}\mathrm{C}_{jsk}\mathrm{C}_{jik} - 8\mathrm{R}_{is}\mathrm{C}_{jsk}\mathrm{C}_{kij} \\
&\quad -\mathrm{R}_{ij}\nabla_k \mathrm{R}\mathrm{C}_{ijk} + \mathrm{R}_{ik}\nabla_j \mathrm{R}\mathrm{C}_{ijk} + 2\mathrm{R}|\mathrm{C}_{ijk}|^2
\end{aligned}$$

$$\begin{aligned}
(\mathrm{R}_{ij}\nabla_l\nabla_l\nabla_k f - \mathrm{R}_{ik}\nabla_l\nabla_l\nabla_j f)\mathrm{C}_{ijk} &= (\mathrm{R}_{ij}\nabla_l(-\mathrm{R}_{lk} + \lambda g_{lk}) - \mathrm{R}_{ik}\nabla_l(-\mathrm{R}_{lj} + \lambda g_{lj}))\mathrm{C}_{ijk} \\
&= -\frac{1}{2}\mathrm{R}_{ij}\nabla_k \mathrm{R}\mathrm{C}_{ijk} + \frac{1}{2}\mathrm{R}_{ik}\nabla_j \mathrm{R}\mathrm{C}_{ijk}
\end{aligned}$$

$$\begin{aligned}
2(\nabla_l \mathrm{R}_{ij}\nabla_l\nabla_k f - \nabla_l \mathrm{R}_{ik}\nabla_l\nabla_j f)\mathrm{C}_{ijk} &= 2((\mathrm{C}_{ijl} + \nabla_j \mathrm{R}_{il} + \frac{1}{4}\nabla_l \mathrm{R} g_{ij} - \frac{1}{4}\nabla_j \mathrm{R} g_{il})(-\mathrm{R}_{lk} + \lambda g_{lk}))\mathrm{C}_{ijk} \\
&\quad -2((\mathrm{C}_{ikl} + \nabla_k \mathrm{R}_{il} + \frac{1}{4}\nabla_l \mathrm{R} g_{ik} - \frac{1}{4}\nabla_k \mathrm{R} g_{il})(-\mathrm{R}_{lj} + \lambda g_{lj}))\mathrm{C}_{ijk} \\
&= -2\mathrm{C}_{ijl}\mathrm{C}_{ijk}\mathrm{R}_{lk} - 2\mathrm{C}_{ijk}\mathrm{R}_{lk}\nabla_j \mathrm{R}_{il} + \frac{1}{2}\mathrm{C}_{ijk}\mathrm{R}_{ik}\nabla_j \mathrm{R} \\
&\quad +2\lambda|\mathrm{C}_{ijk}|^2 + 2\lambda\mathrm{C}_{ijk}\nabla_j \mathrm{R}_{ik} \\
&\quad +2\mathrm{C}_{ikl}\mathrm{C}_{ijk}\mathrm{R}_{lj} + 2\mathrm{C}_{ijk}\mathrm{R}_{lj}\nabla_k \mathrm{R}_{il} - \frac{1}{2}\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k \mathrm{R} \\
&\quad +2\lambda|\mathrm{C}_{ijk}|^2 - 2\lambda\mathrm{C}_{ijk}\nabla_k \mathrm{R}_{ij} \\
&= -2\mathrm{C}_{ilj}\mathrm{C}_{ikj}\mathrm{R}_{lk} - 2\mathrm{C}_{ijk}\mathrm{R}_{lk}\nabla_j \mathrm{R}_{il} + \frac{1}{2}\mathrm{C}_{ijk}\mathrm{R}_{ik}\nabla_j \mathrm{R} + 2\lambda|\mathrm{C}_{ijk}|^2 \\
&\quad -2\lambda(\mathrm{C}_{ijk} + \frac{1}{4}\nabla_k \mathrm{R} g_{ij} - \frac{1}{4}\nabla_j \mathrm{R} g_{ik})\mathrm{C}_{ijk} \\
&\quad -2\mathrm{C}_{ilk}\mathrm{C}_{ijk}\mathrm{R}_{lj} + 2\mathrm{C}_{ijk}\mathrm{R}_{lj}\nabla_k \mathrm{R}_{il} - \frac{1}{2}\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k \mathrm{R} + 2\lambda|\mathrm{C}_{ijk}|^2 \\
&= -2\mathrm{C}_{ilj}\mathrm{C}_{ikj}\mathrm{R}_{lk} - 2\mathrm{C}_{ijk}\mathrm{R}_{lk}\nabla_j \mathrm{R}_{il} + \frac{1}{2}\mathrm{C}_{ijk}\mathrm{R}_{ik}\nabla_j \mathrm{R} + 2\lambda|\mathrm{C}_{ijk}|^2 \\
&\quad -2\mathrm{C}_{ilk}\mathrm{C}_{ijk}\mathrm{R}_{lj} + 2\mathrm{C}_{ijk}\mathrm{R}_{lj}\nabla_k \mathrm{R}_{il} - \frac{1}{2}\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k \mathrm{R}\,.
\end{aligned}$$

Hence, getting back to the main computation and using again the symmetry relations (1.4), we finally get

$$\begin{aligned}
\mathrm{C}_{ijk}\Delta\mathrm{C}_{ijk} &= \frac{1}{2}\nabla_s|\mathrm{C}_{ijk}|^2\nabla_s f + 3\lambda|\mathrm{C}_{ijk}|^2 - \mathrm{R}|\mathrm{C}_{ijk}|^2 \\
&\quad -\frac{3}{2}\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k\mathrm{R} + \frac{3}{2}\mathrm{C}_{ijk}\mathrm{R}_{ik}\nabla_j\mathrm{R} \\
&\quad +4\mathrm{C}_{jsk}\mathrm{C}_{jik}\mathrm{R}_{si} - 8\mathrm{C}_{jsk}\mathrm{C}_{kij}\mathrm{R}_{si} \\
&\quad -2\mathrm{C}_{ijk}\mathrm{R}_{lk}\nabla_j\mathrm{R}_{il} + 2\mathrm{C}_{ijk}\mathrm{R}_{lj}\nabla_k\mathrm{R}_{il} \\
&= \frac{1}{2}\nabla_s|\mathrm{C}_{ijk}|^2\nabla_s f + 3\lambda|\mathrm{C}_{ijk}|^2 - \mathrm{R}|\mathrm{C}_{ijk}|^2 \\
&\quad -3\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k\mathrm{R} + 4\mathrm{C}_{jsk}\mathrm{C}_{jik}\mathrm{R}_{si} - 8\mathrm{C}_{jsk}\mathrm{C}_{kij}\mathrm{R}_{si} - 4\mathrm{C}_{ijk}\mathrm{R}_{lk}\nabla_j\mathrm{R}_{il}
\end{aligned}$$

where in the last passage we applied the skew–symmetry of the Cotton tensor in its last two indexes. The thesis follows. ☐

## 4. THE EVOLUTION EQUATION OF THE COTTON TENSOR IN ANY DIMENSION

In this section we will compute the evolution equation under the Ricci flow of the Cotton tensor $\mathrm{C}_{ijk}$, for every $n$–dimensional Riemannian manifold $(M^n, g(t))$ evolving by Ricci flow.

Among the evolution equations (1.6) we expand the one for the Ricci tensor,

$$\begin{aligned}
\frac{\partial}{\partial t}\mathrm{R}_{ij} = \Delta\mathrm{R}_{ij} &- \frac{2n}{n-2}g^{pq}\mathrm{R}_{ip}\mathrm{R}_{jq} + \frac{2n}{(n-1)(n-2)}\mathrm{R}\mathrm{R}_{ij} + \frac{2}{n-2}|\mathrm{Ric}|^2 g_{ij} \\
&- \frac{2}{(n-1)(n-2)}\mathrm{R}^2 g_{ij} - 2\mathrm{R}^{pq}\mathrm{W}_{pijq}\,.
\end{aligned}$$

Then, we compute the evolution equations of the derivatives of the curvatures assuming, from now on, to be in normal coordinates,

$$\begin{aligned}
\frac{\partial}{\partial t}\nabla_l\mathrm{R} &= \nabla_l\Delta\mathrm{R} + 2\nabla_l|\mathrm{Ric}|^2\,, \\
\frac{\partial}{\partial t}\nabla_s\mathrm{R}_{ij} &= \nabla_s\Delta\mathrm{R}_{ij} - \frac{2n}{n-2}\nabla_s\mathrm{R}_{ip}\mathrm{R}_{jp} - \frac{2n}{n-2}\mathrm{R}_{ip}\nabla_s\mathrm{R}_{jp} + \frac{2n}{(n-1)(n-2)}\nabla_s\mathrm{R}\mathrm{R}_{ij} \\
&\quad + \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_s\mathrm{R}_{ij} + \frac{2}{n-2}\nabla_s|\mathrm{Ric}|^2 g_{ij} - \frac{2}{(n-1)(n-2)}\nabla_s\mathrm{R}^2 g_{ij} \\
&\quad -2\nabla_s\mathrm{R}_{kl}\mathrm{W}_{kijl} - 2\mathrm{R}_{kl}\nabla_s\mathrm{W}_{kijl} + (\nabla_i\mathrm{R}_{sp} + \nabla_s\mathrm{R}_{ip} - \nabla_p\mathrm{R}_{is})\mathrm{R}_{jp} \\
&\quad +(\nabla_j\mathrm{R}_{sp} + \nabla_s\mathrm{R}_{jp} - \nabla_p\mathrm{R}_{js})\mathrm{R}_{ip} \\
&= \nabla_s\Delta\mathrm{R}_{ij} - \frac{n+2}{n-2}\nabla_s\mathrm{R}_{ip}\mathrm{R}_{jp} - \frac{n+2}{n-2}\mathrm{R}_{ip}\nabla_s\mathrm{R}_{jp} + \frac{2n}{(n-1)(n-2)}\nabla_s\mathrm{R}\mathrm{R}_{ij} \\
&\quad + \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_s\mathrm{R}_{ij} + \frac{2}{n-2}\nabla_s|\mathrm{Ric}|^2 g_{ij} - \frac{2}{(n-1)(n-2)}\nabla_s\mathrm{R}^2 g_{ij} \\
&\quad -2\nabla_s\mathrm{R}_{kl}\mathrm{W}_{kijl} - 2\mathrm{R}_{kl}\nabla_s\mathrm{W}_{kijl} + (\nabla_i\mathrm{R}_{sp} - \nabla_p\mathrm{R}_{is})\mathrm{R}_{jp} + (\nabla_j\mathrm{R}_{sp} - \nabla_p\mathrm{R}_{js})\mathrm{R}_{ip} \\
&= \nabla_s\Delta\mathrm{R}_{ij} - \frac{n+2}{n-2}\nabla_s\mathrm{R}_{ip}\mathrm{R}_{jp} - \frac{n+2}{n-2}\mathrm{R}_{ip}\nabla_s\mathrm{R}_{jp} + \frac{2n}{(n-1)(n-2)}\nabla_s\mathrm{R}\mathrm{R}_{ij} \\
&\quad + \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_s\mathrm{R}_{ij} + \frac{2}{n-2}\nabla_s|\mathrm{Ric}|^2 g_{ij} - \frac{2}{(n-1)(n-2)}\nabla_s\mathrm{R}^2 g_{ij} \\
&\quad -2\nabla_s\mathrm{R}_{kl}\mathrm{W}_{kijl} - 2\mathrm{R}_{kl}\nabla_s\mathrm{W}_{kijl} + \mathrm{C}_{spi}\mathrm{R}_{jp} + \mathrm{C}_{spj}\mathrm{R}_{ip} \\
&\quad + \frac{1}{2(n-1)}\mathrm{R}_{jp}[\nabla_i\mathrm{R}g_{sp} - \nabla_p\mathrm{R}g_{is}] + \frac{1}{2(n-1)}\mathrm{R}_{ip}[\nabla_j\mathrm{R}g_{sp} - \nabla_p\mathrm{R}g_{js}]\,,
\end{aligned}$$

where in the last passage we substituted the expression of the Cotton tensor.

We then compute,

$$\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk} &= \frac{\partial}{\partial t}\nabla_k\mathrm{R}_{ij} - \frac{\partial}{\partial t}\nabla_j\mathrm{R}_{ik} - \frac{1}{2(n-1)}\frac{\partial}{\partial t}\big(\nabla_k\mathrm{R}g_{ij} - \nabla_j\mathrm{R}g_{ik}\big)\\
&= \nabla_k\Delta\mathrm{R}_{ij} - \frac{n+2}{n-2}\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp} - \frac{n+2}{n-2}\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp} + \frac{2n}{(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij}\\
&\quad + \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_k\mathrm{R}_{ij} + \frac{2}{n-2}\nabla_k|\mathrm{Ric}|^2 g_{ij} - \frac{2}{(n-1)(n-2)}\nabla_k\mathrm{R}^2 g_{ij}\\
&\quad -2\nabla_k\mathrm{R}_{pl}\mathrm{W}_{pijl} - 2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl} + \mathrm{C}_{kpi}\mathrm{R}_{jp} + \mathrm{C}_{kpj}\mathrm{R}_{ip}\\
&\quad + \frac{\mathrm{R}_{jp}}{2(n-1)}[\nabla_i\mathrm{R}g_{kp} - \nabla_p\mathrm{R}g_{ik}] + \frac{\mathrm{R}_{ip}}{2(n-1)}[\nabla_j\mathrm{R}g_{kp} - \nabla_p\mathrm{R}g_{jk}]\\
&\quad -\nabla_j\Delta\mathrm{R}_{ik} + \frac{n+2}{n-2}\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp} + \frac{n+2}{n-2}\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp} - \frac{2n}{(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&\quad - \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}_{ik} - \frac{2}{n-2}\nabla_j|\mathrm{Ric}|^2 g_{ik} + \frac{2}{n-1}\nabla_j\mathrm{R}^2 g_{ik}\\
&\quad -2\nabla_k\mathrm{R}_{pl}\mathrm{W}_{pijl} - 2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl} - \mathrm{C}_{jpi}\mathrm{R}_{kp} - \mathrm{C}_{jpk}\mathrm{R}_{ip}\\
&\quad - \frac{\mathrm{R}_{kp}}{2(n-1)}[\nabla_i\mathrm{R}g_{jp} - \nabla_p\mathrm{R}g_{ij}] - \frac{\mathrm{R}_{ip}}{2(n-1)}[\nabla_k\mathrm{R}g_{jp} - \nabla_p\mathrm{R}g_{kj}]\\
&\quad + \frac{1}{n-1}\big(\mathrm{R}_{ij}\nabla_k\mathrm{R} - \mathrm{R}_{ik}\nabla_j\mathrm{R}\big)\\
&\quad -\big(\nabla_k\Delta\mathrm{R} + 2\nabla_k|\mathrm{Ric}|^2\big)\frac{g_{ij}}{2(n-1)} + \big(\nabla_j\Delta\mathrm{R} + 2\nabla_j|\mathrm{Ric}|^2\big)\frac{g_{ik}}{2(n-1)}\\
&= \nabla_k\Delta\mathrm{R}_{ij} - \frac{n+2}{n-2}\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp} - \frac{n+2}{n-2}\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp}\\
&\quad + \frac{5n-2}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij} + \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&\quad + \frac{n}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2 g_{ij} - \frac{2}{(n-1)(n-2)}\nabla_k\mathrm{R}^2 g_{ij}\\
&\quad + \mathrm{C}_{kpi}\mathrm{R}_{jp} + \mathrm{C}_{kpj}\mathrm{R}_{ip} - 2\nabla_k\mathrm{R}_{pl}\mathrm{W}_{pijl} - 2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl}\\
&\quad - \frac{1}{2(n-1)}\mathrm{R}_{pj}\nabla_p\mathrm{R}g_{ik}\\
&\quad -\nabla_j\Delta\mathrm{R}_{ik} + \frac{n+2}{n-2}\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp} + \frac{n+2}{n-2}\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp}\\
&\quad - \frac{5n-2}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik} - \frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&\quad - \frac{n}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2 g_{ik} + \frac{2}{(n-1)(n-2)}\nabla_j\mathrm{R}^2 g_{ik}\\
&\quad -\mathrm{C}_{jpi}\mathrm{R}_{kp} - \mathrm{C}_{jpk}\mathrm{R}_{ip} + 2\nabla_j\mathrm{R}_{pl}\mathrm{W}_{pikl} + 2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}\\
&\quad + \frac{1}{2(n-1)}\nabla_l\mathrm{R}\mathrm{R}_{lk}g_{ij} - \frac{1}{2(n-1)}\nabla_k\Delta\mathrm{R}g_{ij} + \frac{1}{2(n-1)}\nabla_j\Delta\mathrm{R}g_{ik}
\end{aligned}$$

and

$$\Delta\mathrm{C}_{ijk} = \Delta\nabla_k\mathrm{R}_{ij} - \Delta\nabla_j\mathrm{R}_{ik} - \frac{1}{2(n-1)}\Delta\nabla_k\mathrm{R}g_{ij} + \frac{1}{2(n-1)}\Delta\nabla_j\mathrm{R}g_{ik}\,,$$

hence,

$$
\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk}-\Delta\mathrm{C}_{ijk} = {} & \nabla_k\Delta\mathrm{R}_{ij}-\nabla_j\Delta\mathrm{R}_{ik}-\Delta\nabla_k\mathrm{R}_{ij}+\Delta\nabla_j\mathrm{R}_{ik}\\
&-\frac{1}{2(n-1)}(\nabla_k\Delta\mathrm{R}g_{ij}-\nabla_j\Delta\mathrm{R}g_{ik}-\Delta\nabla_k\mathrm{R}g_{ij}+\Delta\nabla_j\mathrm{R}g_{ik})\\
&-\frac{n+2}{n-2}(\nabla_k\mathrm{R}_{ip}\mathrm{R}_{jp}+\mathrm{R}_{ip}\nabla_k\mathrm{R}_{jp})+\frac{5n-2}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij}\\
&+\frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&+\frac{n}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2g_{ij}-\frac{2}{(n-1)(n-2)}\nabla_k\mathrm{R}^2g_{ij}\\
&+\mathrm{C}_{kpi}\mathrm{R}_{jp}+\mathrm{C}_{kpj}\mathrm{R}_{ip}-2\nabla_k\mathrm{R}_{pl}\mathrm{W}_{pijl}-2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl}\\
&-\frac{1}{2(n-1)}\mathrm{R}_{jp}\nabla_p\mathrm{R}g_{ik}\\
&+\frac{n+2}{n-2}(\nabla_j\mathrm{R}_{ip}\mathrm{R}_{kp}+\mathrm{R}_{ip}\nabla_j\mathrm{R}_{kp})-\frac{5n-2}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&-\frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&-\frac{n}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2g_{ik}+\frac{2}{(n-1)(n-2)}\nabla_j\mathrm{R}^2g_{ik}\\
&-\mathrm{C}_{jpi}\mathrm{R}_{kp}-\mathrm{C}_{jpk}\mathrm{R}_{ip}+2\nabla_j\mathrm{R}_{pl}\mathrm{W}_{pikl}+2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}\\
&+\frac{1}{2(n-1)}\mathrm{R}_{kp}\nabla_p\mathrm{R}g_{ij}
\end{aligned}
$$

Now to proceed, we need the following commutation rules for the derivatives of the Ricci tensor and of the scalar curvature, where we will employ the decomposition formula of the Riemann tensor (1.1).

$$
\begin{aligned}
\nabla_k\Delta\mathrm{R}_{ij}-\Delta\nabla_k\mathrm{R}_{ij} &= \nabla^3_{kll}\mathrm{R}_{ij}-\nabla^3_{lkl}\mathrm{R}_{ij}+\nabla^3_{lkl}\mathrm{R}_{ij}-\nabla^3_{llk}\mathrm{R}_{ij}\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\mathrm{R}_{klip}\nabla_l\mathrm{R}_{jp}+\mathrm{R}_{kljp}\nabla_l\mathrm{R}_{ip}\\
&\quad +\nabla^3_{lkl}\mathrm{R}_{ij}-\nabla^3_{llk}\mathrm{R}_{ij}\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\frac{1}{2(n-2)}(\mathrm{R}_{ik}\nabla_j\mathrm{R}+\mathrm{R}_{jk}\nabla_i\mathrm{R})\\
&\quad -\frac{1}{n-2}(\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}+\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}-\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}-\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk})\\
&\quad -\frac{1}{n-2}(\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}+\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik})-\frac{1}{2(n-1)(n-2)}(\mathrm{R}\nabla_j\mathrm{R}g_{ik}+\mathrm{R}\nabla_i\mathrm{R}g_{jk})\\
&\quad +\frac{1}{(n-1)(n-2)}(\mathrm{R}\nabla_i\mathrm{R}_{jk}+\mathrm{R}\nabla_j\mathrm{R}_{ik})\\
&\quad +\nabla_l\big(\mathrm{R}_{klip}\mathrm{R}_{pj}+\mathrm{R}_{kljp}\mathrm{R}_{pi}\big)\\
&\quad +\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{ip}+\mathrm{W}_{klip}\nabla_j\mathrm{R}_{jp}\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\frac{1}{2(n-2)}(\mathrm{R}_{ik}\nabla_j\mathrm{R}+\mathrm{R}_{jk}\nabla_i\mathrm{R})\\
&\quad -\frac{1}{n-2}(\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}+\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}-\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}-\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk})\\
&\quad -\frac{1}{n-2}(\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}+\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik})-\frac{1}{2(n-1)(n-2)}(\mathrm{R}\nabla_j\mathrm{R}g_{ik}+\mathrm{R}\nabla_i\mathrm{R}g_{jk})\\
&\quad +\frac{1}{(n-1)(n-2)}(\mathrm{R}\nabla_i\mathrm{R}_{jk}+\mathrm{R}\nabla_j\mathrm{R}_{ik})\\
&\quad +\nabla_l\Big(\frac{1}{n-2}(\mathrm{R}_{ki}g_{pl}\mathrm{R}_{pj}+\mathrm{R}_{pl}g_{ki}\mathrm{R}_{pj}-\mathrm{R}_{li}g_{kp}\mathrm{R}_{pj}-\mathrm{R}_{kp}g_{li}\mathrm{R}_{pj})\\
&\quad -\frac{1}{(n-1)(n-2)}(\mathrm{R}\mathrm{R}_{pj}g_{ki}g_{lp}-\mathrm{R}\mathrm{R}_{pj}g_{kp}g_{il})+\mathrm{W}_{klip}\mathrm{R}_{pj}\\
&\quad +\frac{1}{n-2}(\mathrm{R}_{kj}g_{lp}\mathrm{R}_{pi}+\mathrm{R}_{lp}g_{kj}\mathrm{R}_{pi}-\mathrm{R}_{lj}g_{kp}\mathrm{R}_{pi}-\mathrm{R}_{kp}g_{lj}\mathrm{R}_{pi})\\
&\quad -\frac{1}{(n-1)(n-2)}(\mathrm{R}\mathrm{R}_{pi}g_{kj}g_{pl}-\mathrm{R}\mathrm{R}_{pi}g_{kp}g_{lj})+\mathrm{W}_{kljp}\mathrm{R}_{pi}\Big)\\
&\quad +\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{ip}+\mathrm{W}_{klip}\nabla_j\mathrm{R}_{jp}\\
&= -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\frac{1}{2(n-2)}(\mathrm{R}_{ik}\nabla_j\mathrm{R}+\mathrm{R}_{jk}\nabla_i\mathrm{R})\\
&\quad -\frac{1}{n-2}(\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}+\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}-\mathrm{R}_{lp}\nabla_l\mathrm{R}_{jp}g_{ik}-\mathrm{R}_{lp}\nabla_l\mathrm{R}_{ip}g_{jk})\\
&\quad -\frac{1}{n-2}(\mathrm{R}_{li}\nabla_l\mathrm{R}_{jk}+\mathrm{R}_{lj}\nabla_l\mathrm{R}_{ik})-\frac{1}{2(n-1)(n-2)}(\mathrm{R}\nabla_j\mathrm{R}g_{ik}+\mathrm{R}\nabla_i\mathrm{R}g_{jk})\\
&\quad +\frac{1}{(n-1)(n-2)}(\mathrm{R}\nabla_i\mathrm{R}_{jk}+\mathrm{R}\nabla_j\mathrm{R}_{ik})\\
&\quad +\frac{1}{n-2}(\nabla_p\mathrm{R}_{ki}\mathrm{R}_{pj}+\mathrm{R}_{ki}\nabla_j\mathrm{R}/2+\nabla_p\mathrm{R}g_{ki}\mathrm{R}_{pj}/2\\
&\quad +\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pj}g_{ik}-\nabla_i\mathrm{R}\mathrm{R}_{jk}/2-\mathrm{R}_{pi}\nabla_p\mathrm{R}_{kj}-\nabla_i\mathrm{R}_{kp}\mathrm{R}_{pj}\\
&\quad -\mathrm{R}_{kp}\nabla_i\mathrm{R}_{pj})\\
&\quad -\frac{1}{(n-1)(n-2)}(\nabla_p\mathrm{R}\mathrm{R}_{pj}g_{ik}+\mathrm{R}\nabla_j\mathrm{R}g_{ik}/2-\nabla_i\mathrm{R}\mathrm{R}_{kj}-\mathrm{R}\nabla_i\mathrm{R}_{jk})\\
&\quad +\frac{n-3}{n-2}\mathrm{C}_{kip}\mathrm{R}_{pj}+\mathrm{W}_{klip}\nabla_l\mathrm{R}_{pj}
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{1}{n-2}(\nabla_p\mathrm{R}_{kj}\mathrm{R}_{pi}+\mathrm{R}_{kj}\nabla_i\mathrm{R}/2+\nabla_p\mathrm{R}g_{kj}\mathrm{R}_{pi}/2\\
&+\mathrm{R}_{lp}g_{kj}\nabla_l\mathrm{R}_{pi}-\nabla_j\mathrm{R}\mathrm{R}_{ki}/2-\mathrm{R}_{pj}\nabla_p\mathrm{R}_{ki}-\nabla_j\mathrm{R}_{kp}\mathrm{R}_{pi}-\mathrm{R}_{kp}\nabla_j\mathrm{R}_{pi})\\
&-\frac{1}{(n-1)(n-2)}(\nabla_p\mathrm{R}\mathrm{R}_{pi}g_{kj}+\mathrm{R}\nabla_i\mathrm{R}g_{jk}/2-\nabla_j\mathrm{R}\mathrm{R}_{ki}-\mathrm{R}\nabla_j\mathrm{R}_{ki})\\
&+\frac{n-3}{n-2}\mathrm{C}_{kjp}\mathrm{R}_{pi}+\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{pi}\\
&+\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{ip}+\mathrm{W}_{klip}\nabla_j\mathrm{R}_{jp}\\
=\quad&-\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\frac{n+1}{2(n-1)(n-2)}\mathrm{R}_{kj}\nabla_i\mathrm{R}-\frac{2}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}\\
&+\frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pi}g_{jk}-\frac{1}{n-2}\mathrm{R}_{pj}\nabla_p\mathrm{R}_{ik}-\frac{1}{(n-1)(n-2)}\mathrm{R}\nabla_i\mathrm{R}g_{jk}\\
&+\frac{2}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}_{ik}+\frac{n+1}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ki}-\frac{2}{n-2}\mathrm{R}_{kp}\nabla_i\mathrm{R}_{jp}\\
&+\frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pj}g_{ik}-\frac{1}{n-2}\mathrm{R}_{pi}\nabla_p\mathrm{R}_{jk}-\frac{1}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}g_{ik}\\
&+\frac{2}{(n-1)(n-2)}\mathrm{R}\nabla_i\mathrm{R}_{jk}+2\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{pi}+2\mathrm{W}_{klip}\nabla_l\mathrm{R}_{pj}\\
&+\frac{n-3}{2(n-1)(n-2)}(\nabla_p\mathrm{R}g_{ik}\mathrm{R}_{pj}+\nabla_p\mathrm{R}g_{jk}\mathrm{R}_{pi})\\
&+\frac{n-3}{n-2}(\mathrm{C}_{kip}\mathrm{R}_{pj}+\mathrm{C}_{kjp}\mathrm{R}_{pi})-\frac{1}{n-2}(\nabla_i\mathrm{R}_{kp}\mathrm{R}_{pj}+\nabla_j\mathrm{R}_{kp}\mathrm{R}_{pi})
\end{aligned}
$$

and

$$\nabla_k\Delta\mathrm{R}-\Delta\nabla_k\mathrm{R}=\mathrm{R}_{kllp}\nabla_p\mathrm{R}=-\mathrm{R}_{kp}\nabla_p\mathrm{R}\,.$$

Then, getting back to the main computation, we obtain

$$\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk} - \Delta \mathrm{C}_{ijk} \quad = \quad & -\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij} + \frac{n+1}{2(n-1)(n-2)}\mathrm{R}_{kj}\nabla_i\mathrm{R} \\
& -\frac{2}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip} + \frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pi}g_{jk} \\
& -\frac{1}{n-2}\mathrm{R}_{jp}\nabla_p\mathrm{R}_{ik} - \frac{1}{(n-1)(n-2)}\mathrm{R}\nabla_i\mathrm{R}g_{jk} \\
& +\frac{2}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}_{ik} + \frac{n+1}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ki} \\
& -\frac{2}{n-2}\mathrm{R}_{kp}\nabla_i\mathrm{R}_{pj} + \frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pj}g_{ik} \\
& -\frac{1}{n-2}\mathrm{R}_{pi}\nabla_p\mathrm{R}_{kj} - \frac{1}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}g_{ik} \\
& +\frac{2}{(n-1)(n-2)}\mathrm{R}\nabla_i\mathrm{R}_{jk} + 2\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{pi} + 2\mathrm{W}_{klip}\nabla_l\mathrm{R}_{pj} \\
& +\frac{n-3}{2(n-1)(n-2)}(\nabla_p\mathrm{R}g_{ik}\mathrm{R}_{pj} + \nabla_p\mathrm{R}g_{jk}\mathrm{R}_{pi}) \\
& +\frac{n-3}{n-2}(\mathrm{C}_{kip}\mathrm{R}_{pj} + \mathrm{C}_{kjp}\mathrm{R}_{pi}) \\
& -\frac{1}{n-2}(\nabla_i\mathrm{R}_{kp}\mathrm{R}_{jp} + \nabla_j\mathrm{R}_{kp}\mathrm{R}_{pi}) \\
& +\mathrm{R}_{jp}\nabla_p\mathrm{R}_{ik} - \frac{n+1}{2(n-1)(n-2)}\mathrm{R}_{kj}\nabla_i\mathrm{R} + \frac{2}{n-2}\mathrm{R}_{jp}\nabla_k\mathrm{R}_{ip} \\
& -\frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pi}g_{kj} + \frac{1}{n-2}\mathrm{R}_{pk}\nabla_p\mathrm{R}_{ij} \\
& +\frac{1}{(n-1)(n-2)}\mathrm{R}\nabla_i\mathrm{R}g_{jk} - \frac{2}{(n-1)(n-2)}\mathrm{R}\nabla_k\mathrm{R}_{ij} \\
& -\frac{n+1}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij} + \frac{2}{n-2}\mathrm{R}_{jp}\nabla_i\mathrm{R}_{kp} \\
& -\frac{2}{n-2}\mathrm{R}_{lp}\nabla_p\mathrm{R}_{pk}g_{ij} + \frac{1}{n-2}\mathrm{R}_{pi}\nabla_p\mathrm{R}_{kj} \\
& +\frac{1}{(n-1)(n-2)}\mathrm{R}\nabla_k\mathrm{R}g_{ij} - \frac{2}{(n-1)(n-2)}\mathrm{R}\nabla_i\mathrm{R}_{kj} \\
& -2\mathrm{W}_{jlkp}\nabla_l\mathrm{R}_{pi} - 2\mathrm{W}_{jlip}\nabla_l\mathrm{R}_{pk} \\
& -\frac{n-3}{2(n-2)(n-2)}(\nabla_p\mathrm{R}g_{ij}\mathrm{R}_{pk} + \nabla_p\mathrm{R}g_{jk}\mathrm{R}_{pi}) \\
& -\frac{n-3}{n-2}(\mathrm{C}_{jip}\mathrm{R}_{pk} + \mathrm{C}_{jkp}\mathrm{R}_{pi}) + \frac{1}{n-2}(\nabla_i\mathrm{R}_{pj}\mathrm{R}_{pk} + \nabla_k\mathrm{R}_{jp}\mathrm{R}_{pi}) \\
& +\frac{1}{2(n-1)}(\mathrm{R}_{kp}\nabla_p\mathrm{R}g_{ij} - \mathrm{R}_{jp}\nabla_p\mathrm{R}g_{ki}) - \frac{n+2}{n-2}(\nabla_k\mathrm{R}_{pi}\mathrm{R}_{pj} + \mathrm{R}_{pi}\nabla_k\mathrm{R}_{pj}) \\
& +\frac{n}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2g_{ij} + \frac{5n-2}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij} \\
& +\frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_k\mathrm{R}_{ij} - \frac{2}{(n-1)(n-2)}\nabla_k\mathrm{R}^2g_{ij} \\
& -2\nabla_k\mathrm{R}_{pl}\mathrm{W}_{pijl} - 2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl} \\
& +\mathrm{C}_{kli}\mathrm{R}_{lj} - \frac{1}{2(n-1)}\nabla_l\mathrm{R}\mathrm{R}_{lj}g_{ik} + \mathrm{C}_{klj}\mathrm{R}_{li}
\end{aligned}$$

$$
\begin{aligned}
&+\frac{n+2}{n-2}(\nabla_j\mathrm{R}_{pi}\mathrm{R}_{pk}+\mathrm{R}_{pi}\nabla_j\mathrm{R}_{pk})\\
&-\frac{n}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2 g_{ki}-\frac{5n-2}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&-\frac{2n}{(n-1)(n-2)}\mathrm{R}\nabla_j\mathrm{R}_{ik}+\frac{2}{(n-1)(n-2)}\nabla_j\mathrm{R}^2 g_{ik}\\
&+2\nabla_j\mathrm{R}_{pl}\mathrm{W}_{pikl}+2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}\\
&-\mathrm{C}_{jli}\mathrm{R}_{lk}+\frac{1}{2(n-1)}\nabla_l\mathrm{R}\mathrm{R}_{lk}g_{ij}-\mathrm{C}_{jlk}\mathrm{R}_{li}\\
={}&\frac{1}{n-2}(\mathrm{R}_{pi}\mathrm{C}_{jkp}+\mathrm{R}_{pk}\mathrm{C}_{jip}-\mathrm{C}_{kip}\mathrm{R}_{pj}-\mathrm{C}_{kjp}\mathrm{R}_{pi})\\
&+\Big[\frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pj}+\frac{3}{2(n-1)(n-2)}\nabla_j\mathrm{R}^2\\
&-\frac{1}{2(n-2)}\nabla_p\mathrm{R}\mathrm{R}_{pj}-\frac{n}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2\Big]g_{ik}\\
&-\Big[\frac{2}{n-2}\mathrm{R}_{lp}\nabla_l\mathrm{R}_{pk}+\frac{3}{2(n-1)(n-2)}\nabla_k\mathrm{R}^2\\
&-\frac{1}{2(n-2)}\nabla_p\mathrm{R}\mathrm{R}_{pk}-\frac{n}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2\Big]g_{ij}\\
&-\frac{n-3}{n-2}\mathrm{R}_{kp}\nabla_p\mathrm{R}_{ij}+\frac{n-3}{n-2}\mathrm{R}_{pj}\nabla_p\mathrm{R}_{ik}\\
&+\frac{n}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{pi}+\frac{n+1}{n-2}\nabla_j\mathrm{R}_{pk}\mathrm{R}_{pi}-\frac{2}{n-2}\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&-\frac{4n-3}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}-\frac{1}{n-2}\mathrm{R}_{kp}\nabla_i\mathrm{R}_{pj}+\frac{1}{n-2}\mathrm{R}_{pj}\nabla_i\mathrm{R}_{pk}\\
&-\frac{n}{n-2}\mathrm{R}_{jp}\nabla_k\mathrm{R}_{ip}-\frac{n+1}{n-2}\nabla_k\mathrm{R}_{jp}\mathrm{R}_{ip}+\frac{2}{n-2}\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&+\frac{4n-3}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij}+2\mathrm{W}_{klip}\nabla_l\mathrm{R}_{pj}+2\mathrm{W}_{kljp}\nabla_l\mathrm{R}_{pi}-2\mathrm{W}_{jlkp}\nabla_l\mathrm{R}_{pi}\\
&-2\mathrm{W}_{jlip}\nabla_l\mathrm{R}_{pk}-2\nabla_k\mathrm{R}_{pl}\mathrm{W}_{pijl}-2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl}+2\nabla_j\mathrm{R}_{pl}\mathrm{W}_{pikl}+2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}
\end{aligned}
$$

Now, by means of the very definition of the Cotton tensor (1.2), the identities (1.4), and the symmetries of the Weyl tensor, we substitute

$$
\begin{aligned}
\mathrm{C}_{kpj}-\mathrm{C}_{jpk}={}&-\mathrm{C}_{kjp}-\mathrm{C}_{jpk}=\mathrm{C}_{pkj}\\
\nabla_l\mathrm{R}_{jp}&=\nabla_j\mathrm{R}_{lp}+\mathrm{C}_{pjl}+\frac{1}{2(n-1)}\big(\nabla_l\mathrm{R}g_{pj}-\nabla_j\mathrm{R}g_{pl}\big)\\
\nabla_l\mathrm{R}_{kp}&=\nabla_k\mathrm{R}_{lp}+\mathrm{C}_{pkl}+\frac{1}{2(n-1)}\big(\nabla_l\mathrm{R}g_{pk}-\nabla_k\mathrm{R}g_{pl}\big)\\
\nabla_i\mathrm{R}_{jp}&=\nabla_j\mathrm{R}_{ip}+\mathrm{C}_{pji}+\frac{1}{2(n-1)}\big(\nabla_i\mathrm{R}g_{jp}-\nabla_j\mathrm{R}g_{ip}\big)\\
\nabla_i\mathrm{R}_{kp}&=\nabla_k\mathrm{R}_{ip}+\mathrm{C}_{pki}+\frac{1}{2(n-1)}\big(\nabla_i\mathrm{R}g_{kp}-\nabla_k\mathrm{R}g_{ip}\big)\\
\nabla_p\mathrm{R}_{ij}&=\nabla_j\mathrm{R}_{pi}+\mathrm{C}_{ijp}+\frac{1}{2(n-1)}\big(\nabla_p\mathrm{R}g_{ji}-\nabla_j\mathrm{R}g_{pi}\big)\\
\nabla_p\mathrm{R}_{ik}&=\nabla_k\mathrm{R}_{pi}+\mathrm{C}_{ikp}+\frac{1}{2(n-1)}\big(\nabla_p\mathrm{R}g_{ki}-\nabla_k\mathrm{R}g_{pi}\big)
\end{aligned}
$$

in the last expression above, getting

$$
\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk}-\Delta\mathrm{C}_{ijk} = \ & \frac{1}{n-2}(\mathrm{R}_{pi}\mathrm{C}_{pkj}+\mathrm{R}_{pk}\mathrm{C}_{jip}-\mathrm{C}_{kip}\mathrm{R}_{pj})\\
&+\Big[\frac{2}{n-2}\mathrm{R}_{lp}\Big(\nabla_j\mathrm{R}_{lp}+\mathrm{C}_{pjl}+\frac{1}{2(n-1)}\nabla_l\mathrm{R}g_{pj}\\
&-\frac{1}{2(n-1)}\nabla_j\mathrm{R}g_{pl})\Big)+\frac{3}{2(n-1)(n-2)}\nabla_j\mathrm{R}^2\\
&-\frac{1}{2(n-2)}\nabla_p\mathrm{R}\mathrm{R}_{pj}-\frac{n}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2\Big]g_{ik}\\
&-\Big[\frac{2}{n-2}\mathrm{R}_{lp}\Big(\nabla_k\mathrm{R}_{pl}+\mathrm{C}_{pkl}+\frac{1}{2(n-1)}\nabla_l\mathrm{R}g_{pk}\\
&-\frac{1}{2(n-1)}\nabla_k\mathrm{R}g_{pl}\Big)+\frac{3}{2(n-1)(n-2)}\nabla_k\mathrm{R}^2\\
&-\frac{1}{2(n-2)}\nabla_p\mathrm{R}\mathrm{R}_{pk}-\frac{n}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2\Big]g_{ij}\\
&-\frac{n-3}{n-2}\mathrm{R}_{kp}\Big(\mathrm{C}_{ijp}+\nabla_j\mathrm{R}_{ip}+\frac{1}{2(n-1)}(\nabla_p\mathrm{R}g_{ij}-\nabla_j\mathrm{R}g_{ip})\Big)\\
&+\frac{n-3}{n-2}\mathrm{R}_{pj}\Big(\mathrm{C}_{ikp}+\nabla_k\mathrm{R}_{ip}+\frac{1}{2(n-1)}(\nabla_p\mathrm{R}g_{ik}-\nabla_k\mathrm{R}g_{ip})\Big)\\
&+\frac{n}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{pi}+\frac{n+1}{n-2}\nabla_j\mathrm{R}_{pk}\mathrm{R}_{pi}-\frac{2}{n-2}\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&-\frac{4n-3}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&-\frac{1}{n-2}\mathrm{R}_{kp}\Big(\nabla_j\mathrm{R}_{ip}+\mathrm{C}_{pji}+\frac{1}{2(n-1)}(\nabla_i\mathrm{R}g_{jp}-\nabla_j\mathrm{R}g_{ip})\Big)\\
&+\frac{1}{n-2}\mathrm{R}_{pj}\Big(\nabla_k\mathrm{R}_{ip}+\mathrm{C}_{kpi}+\frac{1}{2(n-1)}(\nabla_i\mathrm{R}g_{kp}-\nabla_k\mathrm{R}g_{ip})\Big)\\
&-\frac{n}{n-2}\mathrm{R}_{jp}\nabla_k\mathrm{R}_{ip}-\frac{n+1}{n-2}\nabla_k\mathrm{R}_{jp}\mathrm{R}_{ip}+\frac{2}{n-2}\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&+\frac{4n-3}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij}\\
&+2\mathrm{C}_{plj}\mathrm{W}_{pikl}-2\mathrm{C}_{plk}\mathrm{W}_{pijl}-2\mathrm{C}_{pil}\mathrm{W}_{jklp}\\
&-2\mathrm{W}_{jklp}\nabla_i\mathrm{R}_{pl}-2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pijl}+2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}\\
= \ & \frac{1}{n-2}\left(\mathrm{R}_{pi}\mathrm{C}_{pkj}+\mathrm{R}_{pk}(\mathrm{C}_{jip}-\mathrm{C}_{pji}-(n-3)\mathrm{C}_{ijp})+\mathrm{R}_{pj}(\mathrm{C}_{pki}-\mathrm{C}_{kip}+(n-3)\mathrm{C}_{ikp})\right)\\
&+\frac{2}{n-2}\mathrm{C}_{pjl}\mathrm{R}_{pl}g_{ik}-\frac{2}{n-2}\mathrm{C}_{pkl}\mathrm{R}_{pl}g_{ij}-2\mathrm{C}_{pjl}\mathrm{W}_{pikl}+2\mathrm{C}_{pkl}\mathrm{W}_{pijl}-2\mathrm{C}_{pil}\mathrm{W}_{jklp}\\
&+g_{ik}\Big[\frac{\nabla_j\mathrm{R}^2}{(n-1)(n-2)}-\frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2\Big]\\
&-g_{ij}\Big[\frac{\nabla_k\mathrm{R}^2}{(n-1)(n-2)}-\frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2\Big]\\
&-\frac{2}{n-2}\mathrm{R}_{jp}\nabla_k\mathrm{R}_{ip}-\frac{n+1}{n-2}\nabla_k\mathrm{R}_{jp}\mathrm{R}_{ip}+\frac{3n-1}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij}+\frac{2}{n-2}\mathrm{R}\nabla_k\mathrm{R}_{ij}\\
&+\frac{2}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip}+\frac{n+1}{n-2}\nabla_j\mathrm{R}_{kp}\mathrm{R}_{ip}-\frac{3n-1}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}-\frac{2}{n-2}\mathrm{R}\nabla_j\mathrm{R}_{ik}\\
&-2\mathrm{W}_{jklp}\nabla_i\mathrm{R}_{lp}-2\mathrm{R}_{lp}\nabla_k\mathrm{W}_{pijl}+2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}\,.
\end{aligned}
$$

then, we substitute again

$$\begin{aligned}
\nabla_k \mathrm{R}_{jp} &= \nabla_p \mathrm{R}_{kj} + \mathrm{C}_{jpk} + \frac{1}{2(n-1)}\big(\nabla_k \mathrm{R} g_{jp} - \nabla_p \mathrm{R} g_{jk}\big)\\
\nabla_j \mathrm{R}_{kp} &= \nabla_p \mathrm{R}_{jk} + \mathrm{C}_{kpj} + \frac{1}{2(n-1)}\big(\nabla_j \mathrm{R} g_{kp} - \nabla_p \mathrm{R} g_{kj}\big)\\
\nabla_k \mathrm{R}_{ij} &= \nabla_i \mathrm{R}_{kj} + \mathrm{C}_{jik} + \frac{1}{2(n-1)}\big(\nabla_k \mathrm{R} g_{ij} - \nabla_i \mathrm{R} g_{jk}\big)\\
\nabla_j \mathrm{R}_{ik} &= \nabla_i \mathrm{R}_{jk} + \mathrm{C}_{kij} + \frac{1}{2(n-1)}\big(\nabla_j \mathrm{R} g_{ik} - \nabla_i \mathrm{R} g_{kj}\big)\,,
\end{aligned}$$

finally obtaining

$$\begin{aligned}
\frac{\partial}{\partial t}\mathrm{C}_{ijk} - \Delta \mathrm{C}_{ijk} \;=\;& \frac{1}{n-2}\left(\mathrm{R}_{pi}\mathrm{C}_{pkj} + \mathrm{R}_{pk}(\mathrm{C}_{jip} - \mathrm{C}_{pji} - (n-3)\mathrm{C}_{ijp}) + \mathrm{R}_{pj}(\mathrm{C}_{pki} - \mathrm{C}_{kip} + (n-3)\mathrm{C}_{ikp})\right)\\
&+\frac{2}{n-2}\mathrm{C}_{pjl}\mathrm{R}_{pl}g_{ik} - \frac{2}{n-2}\mathrm{C}_{pkl}\mathrm{R}_{pl}g_{ij} - 2\mathrm{C}_{pjl}\mathrm{W}_{pikl} + 2\mathrm{C}_{pkl}\mathrm{W}_{pijl} - 2\mathrm{C}_{pil}\mathrm{W}_{jklp}\\
&+g_{ik}\Big[\frac{\nabla_j \mathrm{R}^2}{(n-1)(n-2)} - \frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2\Big]\\
&-g_{ij}\Big[\frac{\nabla_k \mathrm{R}^2}{(n-1)(n-2)} - \frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2\Big]\\
&-\frac{2}{n-2}\mathrm{R}_{jp}\nabla_k\mathrm{R}_{ip} - \frac{n+1}{n-2}\mathrm{R}_{ip}\nabla_p\mathrm{R}_{kj} - \frac{n+1}{n-2}\mathrm{R}_{ip}\mathrm{C}_{jpk}\\
&-\frac{n+1}{2(n-1)(n-2)}\mathrm{R}_{ij}\nabla_k\mathrm{R} + \frac{n+1}{2(n-1)(n-2)}\mathrm{R}_{ip}\nabla_p\mathrm{R} g_{jk}\\
&+\frac{3n-1}{2(n-1)(n-2)}\nabla_k\mathrm{R}\mathrm{R}_{ij} + \frac{2}{n-2}\mathrm{R}(\nabla_i\mathrm{R}_{jk} + \mathrm{C}_{jik} + \frac{1}{2(n-1)}(\nabla_k\mathrm{R} g_{ij} - \nabla_i\mathrm{R} g_{jk}))\\
&+\frac{2}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip} + \frac{n+1}{n-2}\mathrm{R}_{ip}\nabla_p\mathrm{R}_{kj} + \frac{n+1}{n-2}\mathrm{R}_{ip}\mathrm{C}_{kpj} + \frac{n+1}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&-\frac{n+1}{2(n-1)(n-2)}\mathrm{R}_{ip}\nabla_p\mathrm{R} g_{jk} - \frac{3n-1}{2(n-1)(n-2)}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&-\frac{2}{n-2}\mathrm{R}(\nabla_i\mathrm{R}_{jk} + \mathrm{C}_{kij} + \frac{1}{2(n-1)}(\nabla_j\mathrm{R} g_{ik} - \nabla_i\mathrm{R} g_{jk}))\\
&-2\mathrm{W}_{jklp}\nabla_i\mathrm{R}_{lp} - 2\mathrm{R}_{lp}\nabla_k\mathrm{W}_{pijl} + 2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}\\
=\;& \frac{1}{n-2}(\mathrm{R}_{pk}(\mathrm{C}_{jip} - \mathrm{C}_{pji} - (n-3)\mathrm{C}_{ijp}) - \mathrm{R}_{pj}(\mathrm{C}_{kip} - \mathrm{C}_{pki} - (n-3)\mathrm{C}_{ikp})\\
&+(n+2)\mathrm{R}_{pi}\mathrm{C}_{pkj}) + \frac{2}{n-2}(\mathrm{C}_{pjl}\mathrm{R}_{pl}g_{ik} - \mathrm{C}_{pkl}\mathrm{R}_{pl}g_{ij}) + \frac{2}{n-2}\mathrm{R}\mathrm{C}_{ijk}\\
&-2\mathrm{W}_{pikl}\mathrm{C}_{pjl} + 2\mathrm{W}_{pijl}\mathrm{C}_{pkl} - 2\mathrm{C}_{pil}\mathrm{W}_{jklp}\\
&+g_{ik}\Big[\frac{\nabla_j \mathrm{R}^2}{2(n-1)(n-2)} - \frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2\Big]\\
&-g_{ij}\Big[\frac{\nabla_k \mathrm{R}^2}{2(n-1)(n-2)} - \frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2\Big]\\
&-\frac{2}{n-2}\mathrm{R}_{jp}\nabla_k\mathrm{R}_{ip} + \frac{1}{n-2}\nabla_k\mathrm{R}\mathrm{R}_{ij}\\
&+\frac{2}{n-2}\mathrm{R}_{kp}\nabla_j\mathrm{R}_{ip} - \frac{1}{n-2}\nabla_j\mathrm{R}\mathrm{R}_{ik}\\
&+2\mathrm{R}_{lp}\nabla_j\mathrm{W}_{pikl} - 2\mathrm{R}_{lp}\nabla_k\mathrm{W}_{pijl}\,,
\end{aligned}$$

where in the last passage we used again the identities (1.4) and the fact that

$$\mathrm{W}_{jklp}\nabla_i\mathrm{R}_{lp} = \mathrm{W}_{jkpl}\nabla_i\mathrm{R}_{pl} = \mathrm{W}_{jkpl}\nabla_i\mathrm{R}_{lp} = -\mathrm{W}_{jklp}\nabla_i\mathrm{R}_{lp}\,.$$

Hence, we can resume this long computation in the following proposition, getting back to a generic coordinate basis.

**Proposition 4.1.** *During the Ricci flow of a $n$–dimensional Riemannian manifold $(M^n, g(t))$, the Cotton tensor satisfies the following evolution equation*

$$\begin{aligned}
\big(\partial_t - \Delta\big)\mathrm{C}_{ijk} \;=\;& \frac{1}{n-2}\,[g^{pq}\mathrm{R}_{pj}(\mathrm{C}_{kqi} + \mathrm{C}_{qki} + (n-3)\mathrm{C}_{ikp})\\
&+(n+2)g^{pq}\mathrm{R}_{ip}\mathrm{C}_{qkj} - g^{pq}\mathrm{R}_{pk}(\mathrm{C}_{jqi} + \mathrm{C}_{qji} + (n-3)\mathrm{C}_{ijq})]\\
&+\frac{2}{n-2}\mathrm{R}\mathrm{C}_{ijk} + \frac{2}{n-2}\mathrm{R}^{ql}\mathrm{C}_{qjl}g_{ik} - \frac{2}{n-2}\mathrm{R}^{ql}\mathrm{C}_{qkl}g_{ij}\\
&+\frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2 g_{ij} - \frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2 g_{ik}\\
&+\frac{\mathrm{R}}{(n-1)(n-2)}\nabla_j\mathrm{R}g_{ik} - \frac{\mathrm{R}}{(n-1)(n-2)}\nabla_k\mathrm{R}g_{ij}\\
&+\frac{2}{n-2}g^{pq}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{qi} - \frac{2}{n-2}g^{pq}\mathrm{R}_{pj}\nabla_k\mathrm{R}_{qi} + \frac{1}{n-2}\mathrm{R}_{ij}\nabla_k\mathrm{R} - \frac{1}{n-2}\mathrm{R}_{ik}\nabla_j\mathrm{R}\\
&-2g^{pq}\mathrm{W}_{pikl}\mathrm{C}_{qjl} + 2g^{pq}\mathrm{W}_{pijl}\mathrm{C}_{qkl} - 2g^{pq}\mathrm{W}_{jklp}\mathrm{C}_{qil} + 2g^{pq}\mathrm{R}_{pl}\nabla_j\mathrm{W}_{qikl} - 2g^{pq}\mathrm{R}_{pl}\nabla_k\mathrm{W}_{qijl}\,.
\end{aligned}$$

*In particular if the Cotton tensor vanishes identically along the flow we obtain,*

$$\begin{aligned}
0 \;=\;& \frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2 g_{ij} - \frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2 g_{ik}\\
&+\frac{\mathrm{R}}{(n-1)(n-2)}\nabla_j\mathrm{R}g_{ik} - \frac{\mathrm{R}}{(n-1)(n-2)}\nabla_k\mathrm{R}g_{ij}\\
&+\frac{2}{n-2}g^{pq}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{qi} - \frac{2}{n-2}g^{pq}\mathrm{R}_{pj}\nabla_k\mathrm{R}_{qi} + \frac{1}{n-2}\mathrm{R}_{ij}\nabla_k\mathrm{R} - \frac{1}{n-2}\mathrm{R}_{ik}\nabla_j\mathrm{R}\\
&-2g^{pq}\mathrm{W}_{jklp}\mathrm{C}_{qil} + 2g^{pq}\mathrm{R}_{pl}\nabla_j\mathrm{W}_{qikl} - 2g^{pq}\mathrm{R}_{pl}\nabla_k\mathrm{W}_{qijl}\,,
\end{aligned}$$

*while, in virtue of* (1.3)*, if the Weyl tensor vanishes along the flow we obtain*

$$\begin{aligned}
0 \;=\;& \frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2 g_{ij} - \frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2 g_{ik}\\
&+\frac{\mathrm{R}}{(n-1)(n-2)}\nabla_j\mathrm{R}g_{ik} - \frac{\mathrm{R}}{(n-1)(n-2)}\nabla_k\mathrm{R}g_{ij}\\
&+\frac{2}{n-2}g^{pq}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{qi} - \frac{2}{n-2}g^{pq}\mathrm{R}_{pj}\nabla_k\mathrm{R}_{qi} + \frac{1}{n-2}\mathrm{R}_{ij}\nabla_k\mathrm{R} - \frac{1}{n-2}\mathrm{R}_{ik}\nabla_j\mathrm{R}\,.
\end{aligned}$$

**Corollary 4.2.** *During the Ricci flow of a $n$–dimensional Riemannian manifold $(M^n, g(t))$, the squared norm of the Cotton tensor satisfies the following evolution equation, in an orthonormal basis,*

$$\begin{aligned}
\big(\partial_t - \Delta\big)|\mathrm{C}_{ijk}|^2 \;=\;& -2|\nabla\mathrm{C}_{ijk}|^2 - \frac{8n}{n-2}\mathrm{C}_{ipk}\mathrm{C}_{iqk}\mathrm{R}_{pq} + \frac{4n+16}{n-2}\mathrm{C}_{ipk}\mathrm{C}_{kqi}\mathrm{R}_{pq}\\
&+\frac{4}{n-2}\mathrm{R}|\mathrm{C}_{ijk}|^2 + \frac{8}{n-2}\mathrm{C}_{ijk}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{pi} + \frac{4}{n-2}\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k\mathrm{R}\\
&+8\mathrm{C}_{ijk}\mathrm{R}_{lp}\nabla_j\mathrm{W}_{pikl} - 8\mathrm{C}_{ijk}\mathrm{C}_{pjl}\mathrm{W}_{pikl} - 4\mathrm{C}_{jpi}\mathrm{C}_{ljk}\mathrm{W}_{pikl}\,.
\end{aligned}$$

$$\begin{aligned}
(\partial_t-\Delta)|\mathrm{C}_{ijk}|^2 \;=\;& -2|\nabla\mathrm{C}_{ijk}|^2+2\mathrm{C}_{ijk}\Big[\frac{1}{n-2}\,[(\mathrm{R}_{pj}(\mathrm{C}_{kpi}+\mathrm{C}_{pki}+(n-3)\mathrm{C}_{ikp})\\
&+(n+2)\mathrm{R}_{pi}\mathrm{C}_{pkj}-\mathrm{R}_{pk}(\mathrm{C}_{jpi}+\mathrm{C}_{pji}+(n-3)\mathrm{C}_{ijp}))\\
&+\frac{2}{n-2}\mathrm{R}\mathrm{C}_{ijk}+\frac{2}{n-2}\mathrm{R}_{ql}\mathrm{C}_{qjl}g_{ik}-\frac{2}{n-2}\mathrm{R}_{ql}\mathrm{C}_{qkl}g_{ij}\\
&+\frac{1}{(n-1)(n-2)}\nabla_k|\mathrm{Ric}|^2g_{ij}-\frac{1}{(n-1)(n-2)}\nabla_j|\mathrm{Ric}|^2g_{ik}\\
&+\frac{\mathrm{R}}{(n-1)(n-2)}\nabla_j\mathrm{R}g_{ik}-\frac{\mathrm{R}}{(n-1)(n-2)}\nabla_k\mathrm{R}g_{ij}\\
&+\frac{2}{n-2}\mathrm{R}_{qk}\nabla_j\mathrm{R}_{qi}-\frac{2}{n-2}\mathrm{R}_{qj}\nabla_k\mathrm{R}_{qi}+\frac{1}{n-2}\mathrm{R}_{ij}\nabla_k\mathrm{R}-\frac{1}{n-2}\mathrm{R}_{ik}\nabla_j\mathrm{R}\\
&-2\mathrm{W}_{pikl}\mathrm{C}_{pjl}+2\mathrm{W}_{pijl}\mathrm{C}_{pkl}-2\mathrm{W}_{jklp}\mathrm{C}_{pil}+2\mathrm{R}_{pl}\nabla_j\mathrm{W}_{pikl}-2\mathrm{R}_{pl}\nabla_k\mathrm{W}_{pikl}\Big]\\
=\;& -2|\nabla\mathrm{C}_{ijk}|^2-\frac{8n}{n-2}\mathrm{C}_{ipk}\mathrm{C}_{iqk}\mathrm{R}_{pq}+\frac{4n+16}{n-2}\mathrm{C}_{ipk}\mathrm{C}_{kqi}\mathrm{R}_{pq}\\
&+\frac{4}{n-2}\mathrm{R}|\mathrm{C}_{ijk}|^2+\frac{8}{n-2}\mathrm{C}_{ijk}\mathrm{R}_{pk}\nabla_j\mathrm{R}_{pi}+\frac{4}{n-2}\mathrm{C}_{ijk}\mathrm{R}_{ij}\nabla_k\mathrm{R}\\
&+8\mathrm{C}_{ijk}\mathrm{R}_{lp}\nabla_j\mathrm{W}_{pikl}-8\mathrm{C}_{ijk}\mathrm{C}_{pjl}\mathrm{W}_{pikl}-4\mathrm{C}_{jpi}\mathrm{C}_{ljk}\mathrm{W}_{pikl}\,.
\end{aligned}$$

*Remark* 4.3. Notice that if $n=3$ the two formulas in Proposition 4.1 and Corollary 4.2 become the ones in Proposition 2.1 and Corollary 2.5.

## REFERENCES

1. A. L. Besse, *Einstein manifolds*, Springer–Verlag, Berlin, 2008.
2. H.-D. Cao, B.-L. Chen, and X.-P. Zhu, *Recent developments on Hamilton's Ricci flow*, Surveys in differential geometry. Vol. XII. Geometric flows, vol. 12, Int. Press, Somerville, MA, 2008, pp. 47–112.
3. H.-D. Cao and Q. Chen, *On locally conformally flat gradient steady Ricci solitons*, Trans. Amer. Math. Soc. **364** (2012), 2377–2391.
4. G. Catino and C. Mantegazza, *Evolution of the Weyl tensor under the Ricci flow*, Ann. Inst. Fourier (2011), 1407–1435.
5. S. Gallot, D. Hulin, and J. Lafontaine, *Riemannian Geometry*, Springer–Verlag, 1990.
6. R. S. Hamilton, *Three–manifolds with positive Ricci curvature*, J. Diff. Geom. **17** (1982), no. 2, 255–306.

(Carlo Mantegazza) SCUOLA NORMALE SUPERIORE DI PISA, PIAZZA CAVALIERI 7, PISA, ITALY, 56126
*E-mail address*, C. Mantegazza: `c.mantegazza@sns.it`

(Carlo Mantegazza) SCUOLA NORMALE SUPERIORE DI PISA, PIAZZA CAVALIERI 7, PISA, ITALY, 56126
*E-mail address*, S. Mongodi: `samuele.mongodi@gmail.com`

(Michele Rimoldi) DIPARTIMENTO DI SCIENZA E ALTA TECNOLOGIA, UNIVERSITÁ DEGLI STUDI DELL'INSUBRIA, VIA VALLEGGIO 11, COMO, ITALY, 22100
*E-mail address*, M. Rimoldi: `michele.rimoldi@gmail.com`